

Longstanding Exercise Therapy in Patients with Rheumatoid Arthritis

No registrations found.

Ethical review	Positive opinion
Status	Recruiting
Health condition type	-
Study type	Interventional

Summary

ID

NL-OMON19895

Source

NTR

Brief title

L-EXTRA study

Health condition

Rheumatoid Arthritis

Sponsors and support

Primary sponsor: Leiden University Medical Centre, LUMC

Source(s) of monetary or material Support: Ministry of Health, Welfare and Sport and ReumaNederland and Koninklijk Nederlands Genootschap voor Fysiotherapie

Intervention

Outcome measures

Primary outcome

The primary outcome measure of effectiveness is the individual level of functioning (activities and participation), as measured with the Patient-Specific Complaints instrument (PSC) at 52 weeks.

Secondary outcome

Secondary outcome measures include the 10-item Patient Reported Outcomes Measurement Information System (PROMIS), the Health Assessment Questionnaire-Disability Index (HAQ-DI) and a 6 minute walk test for functional ability; and the RAQoL and SF-36 for health related Quality of life. In order to address the topic of cost-effectiveness, the EuroQoL (EQ-5D-5L) for health valuation and comprehensive measurements of costs for an economic analysis will be administered as well.

Study description

Background summary

Rationale:

Rheumatoid arthritis (RA) is a chronic systemic disease, mainly characterized by arthritis of the peripheral joints, affecting about 0.4-1% of the population, with women being more often affected than men. During the course of their disease, most patients use short, intermittent physical therapy treatment (active exercise therapy) in order to prevent or diminish limitations in activities or participation. However, there is a substantial subgroup of patients with RA (< 5%) with joint damage or persistent high disease activity, complications of the disease or its treatment or comorbidity, resulting in complex limitations in activities and participation. Due to the individual nature of the problems and resulting disability, in this group the exercise therapy treatment is currently highly personalized and usually longstanding (i.e. longer than 12 weeks and more than 20 sessions per year). Only one high quality randomized controlled trial proved the effectiveness and safety of longstanding exercise therapy, however this concerned a selected group of patients with stable RA and little joint damage (De Jong 2003, 2004). Research on effectiveness of longstanding exercise therapy in the abovementioned patient group with complex limitations in activities and participation is absent.

Objectives:

This study aims to underpin the delivery of longstanding exercise therapy in the subgroup of patients with RA and complex disability.

There are 3 research questions to be addressed:

1. Is longstanding, optimized active exercise therapy more effective with respect to functional ability than usual care in patients with RA with severe functional disability over a period of 52 weeks?
2. Which option (longstanding, optimized active exercise therapy or usual care) is more cost-effective?
3. What is the long-term course of functional ability, health status and health care consumption of patients using longstanding, optimized active exercise therapy?

Study design:

Randomized, controlled trial comparing longstanding, active exercise therapy with usual care

(1:1). After the experimental period of 52 weeks at which the primary end-point is assessed, the intervention will be continued in the intervention group. Effects will be monitored at follow-up measurements after 104 weeks and after 156 and 208 weeks or at the end of the study, depending on date of inclusion. At 52 weeks, the intervention will also be offered to the patients allocated to the usual care group.

Study population:

215 patients with a confirmed diagnosis of RA, with persistent high disease activity, joint damage, complications of the disease or its treatment or comorbidity, resulting in complex limitations in activities and participation.

Intervention and control conditions:

The intervention concerns longstanding, intensive active exercise therapy (52 weeks), aimed at the improvement of specific individual limitations in daily activities and participation. It consists of a standardized program comprising active modalities (functional exercises, aerobic exercises, muscle strengthening and flexibility/joint range of motion exercises), with the type of exercises, their intensity, frequency, duration, site of delivery (practice or at home) and progression being tailored to the individual patients' functional disability and ensuing needs and goals. The control condition consists of care as usual, left to the discretion of the treating physicians and the patients.

Main study parameters/endpoints:

Assessments are done at 0, 12, 26, 52, 104, and 156 and 208 weeks or end of the study (48 months). The primary outcome measure of effectiveness is the individual level of functioning (activities and participation), as measured with the Patient-Specific Complaints instrument (PSC) at 52 weeks.

Secondary outcome measures include the 10-item Patient Reported Outcomes Measurement Information System (PROMIS), the Health Assessment Questionnaire-Disability Index (HAQ-DI) and a 6 minute walk test for functional ability; and the RAQoL and SF-36 for health related Quality of life. In order to address the topic of cost-effectiveness, the EuroQoL (EQ-5D-5L) for health valuation and comprehensive measurements of costs for an economic analysis will be administered as well.

Apart from the primary and secondary outcome measures, sociodemographic and disease characteristics, will be recorded. The presence of comorbidity will be recorded by means of the comorbidity questionnaire developed by the Dutch Central Bureau of Statistics.

In addition, an anchor question regarding the perceived effectiveness will be added in all cases where longstanding exercise therapy was used (intervention and control group if starting after 52 weeks) ("has the exercise therapy changed your daily functioning"), as well as a short questionnaire on patient satisfaction with treatment. Furthermore, we will evaluate the content of care provided and the compliance of the patients in the intervention group by asking the patients to fill out a registration form on the frequency, duration, and content of treatment. In a random sample of 10% we will validate the registration form with the records of the physical therapist (content, duration). The perception of any side effects of exercise therapy will be recorded. If treatment is discontinued, the reasons will be recorded.

Nature and extent of the burden and risks associated with participation, benefit and group relatedness:

The intervention concerns exercise therapy delivered by trained primary care physical therapists according to a standardized protocol, with no extra risks as compared to the regular delivery of primary care exercise therapy. Regarding the burden, the assessments mainly consist of the completion of questionnaires at home (max. 1 hour) and a site visit with one performance test (6-minute walk test) and the answering of a number of questions (max. total visit duration ½ hour). The maximum number of site visits is 5 (1 screening and maximum 4 evaluation visits at: baseline, 26 weeks, 52 weeks (primary endpoint) and 104 weeks. Evaluations at 156 and 208 weeks/end of study only consist of questionnaires.

Study objective

This study aims to underpin the delivery of longstanding exercise therapy in the subgroup of patients with RA and complex disability.

There are 3 research questions to be addressed:

1. Is longstanding, optimized active exercise therapy more effective with respect to functional ability than usual care in patients with RA with severe functional disability over a period of 52 weeks?
2. Which option (longstanding, optimized active exercise therapy or usual care) is more cost-effective?
3. What is the long-term course of functional ability, health status and health care consumption of patients using longstanding, optimized active exercise therapy?

Study design

Assessments are done at 0, 12, 26, 52, 104, and 156 and 208 weeks or end of the study (48 months). The primary outcome measure of effectiveness is the individual level of functioning (activities and participation), as measured with the Patient-Specific Complaints instrument (PSC) at 52 weeks.

Intervention

Longstanding optimized active exercise therapy (intervention) or usual care (control)

Contacts

Public

LUMC, Leiden
Thea Vliet Vlieland

071-5263613

Scientific

LUMC, Leiden
Thea Vliet Vlieland

Eligibility criteria

Inclusion criteria

A confirmed diagnosis of RA, with persistent high disease activity, joint damage, complications of the disease or its treatment or comorbidity, resulting in complex limitations in activities and participation.

Exclusion criteria

Recent (<3 months) or present individual physiotherapy or exercise therapy or multidisciplinary rehabilitation treatment

Study design

Design

Study type:	Interventional
Intervention model:	Parallel
Allocation:	Randomized controlled trial
Masking:	Single blinded (masking used)
Control:	Active

Recruitment

NL	
Recruitment status:	Recruiting
Start date (anticipated):	01-01-2020
Enrollment:	215
Type:	Anticipated

IPD sharing statement

Plan to share IPD: Undecided

Ethics review

Positive opinion

Date: 13-12-2019

Application type: First submission

Study registrations

Followed up by the following (possibly more current) registration

ID: 52885

Bron: ToetsingOnline

Titel:

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register	ID
NTR-new	NL8235
CCMO	NL69866.058.19
OMON	NL-OMON52885

Study results