

# Deployment of dedicated nursing staff to stimulate use of clozapine.

No registrations found.

<b>Ethical review</b>	Positive opinion
<b>Status</b>	Pending
<b>Health condition type</b>	-
<b>Study type</b>	Interventional

## Summary

### ID

NL-OMON20244

### Source

Nationaal Trial Register

### Brief title

CLOZ-NP

### Health condition

Non-affective psychotic disorder.

Niet-affectieve psychotische stoornis.

Titel: Deployment of dedicated nursing staff to stimulate use of clozapine.

Hypothese: Physicians in teams of Type/Condition A (where they can delegate clozapine monitoring tasks to a Nurse Practitioner) will prescribe clozapine more often than physicians in teams of Type/Condition B (where they should performing monitoring tasks themselves).

Blinding: Single-blind

Primaire uitkomst: Proportion of patients indicated for clozapine (in condition A vs. B) who actually start with this drug.

Bij korte samenvatting: Primary objective: to examine whether a physician will treat more patients (who have an indication for clozapine) with clozapine, if he can delegate weekly monitoring tasks and discussion of common side-effects to a Nurse Practitioner (NP).

## Sponsors and support

**Primary sponsor:** Rivierduinen, GGZ Leiden, the Netherlands and Maastricht University, the Netherlands.

**Source(s) of monetary or material Support:** GGZ Leiden, Rivierduinen, the Netherlands

## Intervention

## Outcome measures

### Primary outcome

- 1a. Proportion of missed weekly lab exams (white blood cells) in condition A vs. B.
- 1b. Number of days the lab exams (white blood cells) are delayed, divided by total number of required lab exams in condition A vs. B.

### Secondary outcome

1. Duration of clozapine use in condition A vs. B.
2. Clinical Global Impression-Schizophrenia scale (CGI-SCH) score of change in condition A vs. B.
3. Proportion of patients indicated for clozapine (in condition A vs. B) who actually start with this drug.
4. Proportion of patients for whom clozapine is indicated.
5. Qualitative data to explore how potentially hazardous side-effects were handled.

## Study description

### Background summary

Background: The antipsychotic drug clozapine is more effective than other antipsychotics. However, it remains underused, most likely because it is more labour-intensive as white blood cells should be monitored weekly during the first 18 weeks. Especially for outpatients,

this generates a lot of extra work for the physician (psychiatrist/resident). A recent survey concluded that professionals “perceived the presence of dedicated staff to arrange and monitor the initiation of clozapine in outpatients as the factor that would enable the use of clozapine most”. The exact proportion of patients with Non-Affective Psychotic Disorder (NAPD) in an outpatient setting who should be treated with clozapine is uncertain.

Primary objective: To demonstrate that delegation of clozapine monitoring tasks to a NP is at least as safe as Treatment As Usual (TAU).

Secondary objectives: 1. To examine whether collaboration between a physician and a NP is associated with a longer retention of patients on clozapine. 2. To examine whether collaboration between a physician and a NP is associated with a better outcome in patients who start with clozapine. 3. To examine whether a physician will treat more patients (who have an indication for clozapine) with this drug if he can delegate weekly monitoring tasks and discussion of common side-effects to a Nurse Practitioner (NP). 4. To estimate the proportion of outpatients with NAPD for whom clozapine is indicated, by type of indication 5. To examine the difference between condition A and B in promptness and accuracy of treatment of potentially hazardous side-effects.

In this study we will randomize at least 16 teams into 2 conditions: (A) Collaboration in clozapine treatment between physician and NP, versus (B) TAU: physician performs clozapine monitoring himself.

## **Study objective**

Delegation of clozapine monitoring tasks to a Nurse Practitioner (condition A) is at least as safe as Treatment As Usual: physicians perform monitoring tasks themselves (condition B).

## **Study design**

04-06-2015: Education of physicians and NPs: during one afternoon psychiatrists of the Dutch Clozapine Collaboration Group ([www.clozapinepluswerkgroep.nl](http://www.clozapinepluswerkgroep.nl)) educate physicians and NPs of all participating teams about the diagnosis of treatment-resistance and about other indications for clozapine.

Between 04-06-2015 and 01-10-2015: Baseline assessment: The independent researcher, the responsible physician and NP of each team discuss every patient with NAPD to determine if the patient: a. already uses clozapine; b. has used clozapine and has discontinued this drug (justified or unjustified); c. has an indication for clozapine, with information on type(s) of indication; d. has no indication for clozapine.

September 2015: Training of nurse practitioners.

Between 01-10-2015 and 04-02-2017: One-year study period during which eligible patients will start using clozapine or not. Of all patients that start using clozapine, data will be collected during 18 weeks. Since a patient started in the last week of this year should be monitored for 18 weeks, the study will last an extra 4 months (until February 2017).

Sept 2016-jan. 2017: an extra Treatment As Usual group will be created. This group consists of patients from the intervention teams, who have started clozapine before the start of the

trial.

## **Intervention**

1. Education of physicians and NPs about indications for clozapine.
2. Physicians and NPs screen all patients in their team for a possible indication for clozapine.
3. Randomization of teams into condition A or B (treatment as usual).
4. Training of NPs from teams in condition A in management of patients on clozapine.
5. One-year study period will be study period of 15 months

## **Contacts**

### **Public**

GGZ Leiden

Yvonne van der Zalm  
Sandifortdreef 19

Leiden 2333 ZZ  
The Netherlands  
06-11519672

### **Scientific**

GGZ Leiden

Yvonne van der Zalm  
Sandifortdreef 19

Leiden 2333 ZZ  
The Netherlands  
06-11519672

## **Eligibility criteria**

### **Inclusion criteria**

Teams:

- Outpatient teams in which a nurse practitioner participates

Patients:

- Age 18-55
- Diagnosis: non-affective psychotic disorder

## Exclusion criteria

NA

## Study design

### Design

Study type:	Interventional
Intervention model:	Parallel
Allocation:	Randomized controlled trial
Masking:	Single blinded (masking used)
Control:	Active

### Recruitment

NL	
Recruitment status:	Pending
Start date (anticipated):	04-06-2015
Enrollment:	16
Type:	Anticipated

### IPD sharing statement

**Plan to share IPD:** Undecided

## Ethics review

Positive opinion	
Date:	28-04-2015
Application type:	First submission

## Study registrations

### Followed up by the following (possibly more current) registration

No registrations found.

### Other (possibly less up-to-date) registrations in this register

No registrations found.

### In other registers

Register	ID
NTR-new	NL4988
NTR-old	NTR5135
Other	- : RTP0 937a

## Study results

### Summary results

None yet