AVARIJN2 study

No registrations found.

Ethical review Not applicable

Status Pending

Health condition type -

Study type Observational non invasive

Summary

ID

NL-OMON21621

Source

NTR

Brief title

AVARIJN2 (Aortic VAlve RIJNMOND 2)

Health condition

Aortic valve stenosis

Sponsors and support

Primary sponsor: Erasmus Medical Center, Department of Cardio-thoracic Surgery **Source(s) of monetary or material Support:** Erasmus Medical Center, Department of Cardio-thoracic Surgery

Intervention

Outcome measures

Primary outcome

The primary objective is to determine the quality of life assessed with the SF-36V2¢â questionnaire in asymptomatic patients with severe aortic stenosis, the mortality (all-cause and specific cause) as well as the determination of the aortic valve function by an echocardiogram.

Secondary outcome

The secondary objective is to depict the emanated complications occurred during follow-up.

Study description

Background summary

The European as well as the American guidelines recommend aortic valve replacement (AVR) for patients with severe aortic stenosis (AS) having the concurrent symptoms related to AS (1, 2). Undergoing AVR for severe aortic stenosis results in better outcome compared to patients remaining asymptomatic and therefore not undergoing AVR (3). An overall improvement in functional quality of life, in patients undergoing AVR, has also been noted (4).

However, data focusing on asymptomatic patients is relative scarce. Recent studies did compare the operative and long-term cardiac mortality, as well as different parameters in asymptomatic (NYHA I) patients with severe aortic stenosis undergoing initial surgery versus a conservative treatment strategy (5, 6). Moreover, a randomized controlled trial, the AVATAR study, has begun recruiting patients, wherein they will be comparing an early surgery strategy with a watchful waiting strategy (7). The primary outcome measure, as described in published rationale and design paper of the AVATAR study, is a global composite of 'hard' endpoints including the all-cause mortality (7).

Whereas this study will provide new insight pertaining 'hard' endpoints as the mortality, quality of life assessment in a specific population has hardly been quantified, wherefore we want to conduct this study. To provide insight in the long-term follow-up focusing on the quality of life of patients with asymptomatic severe AS, also to depict the mortality and to evaluate the aortic valve function as obtained with an echocardiogram during the follow-up, to fully understand the effect severe aortic stenosis has in the asymptomatic population.

Hypothesis

We hypothesize that a significant proportion of patients with severe aortic stenosis is not referred for surgical treatment because of co-morbidities and/or old age, or is operated late in the course of the disease due to underestimation of disease severity and progression. We also hypothesize to see altered echocardiographic parameters in these asymptomatic severe AS patients.

Study objectives

Primary Objective(s): The primary objective is to determine the quality of life assessed with the SF-36V2 questionnaire in asymptomatic patients with severe aortic stenosis, the mortality (all-cause and specific cause) as well as the determination of the aortic valve function by an echocardiogram.

Secondary Objective(s): The secondary objective is to depict the emanated complications occurred during follow-up.

Study design

It is a hospital-based cohort study. All asymptomatic patients with severe aortic stenosis who participated in the AVARIJN-study will be selected. All patients who survived will be invited by telephone and letter to visit the outpatient clinic to undergo an echocardiogram and to fill the SF-36V2¢â Health Survey questionnaire. Further data will be obtained by the use of medical files and the hospital information system.

Methods

The asymptomatic patients included in the AVARIJN study will be provided with study information, and invited to the Erasmus Medical Center for participation in this study. Written informed consent will be obtained prior to participation. Echocardiographic inclusion criteria in the AVARIJN study was at least one of the following: (1) aortic valve area ≤ 1 cm2, (2) maximal trans aortic jet velocity ≥ 4 m/s, (3) peak aortic gradient ≥ 64 mmHg and (4) aortic valve / left ventricular outflow tract velocity time integral ratio ≥ 4 .

Patient characteristics, functional status, quality of life and treatment undergone will be sought retrospectively. The patients will undergo an extra echocardiogram and have to fill in the SF-36V2 Health Survey questionnaire. We expect to include around 45 patients, as the study population of the AVARIJN study consists of 59 asymptomatic patients, with the expectation being to include between 85-95% of the eligible patients.

Study objective

We hypothesize that, asymptomatic patients with severe aortic stenosis, receiving an aortic valve replacement have a better quality of life compared to patients without aortic valve replacement.

Study design

Accrual time: 01-07-2016 till 31-12-2017

Study end: 31-12-2018

Intervention

Retrospective follow-up of the asymptomatic patients in the AVARIJN study

Contacts

Public

Erasmus MC Rotterdam
Dept. Cardio Thoracic Surgery
's Gravendijkwal 230
A.P. Kappetein
Rotterdam 3015 CE
The Netherlands
+31 10 7032150

Scientific

Erasmus MC Rotterdam
Dept. Cardio Thoracic Surgery
's Gravendijkwal 230
A.P. Kappetein
Rotterdam 3015 CE
The Netherlands
+31 10 7032150

Eligibility criteria

Inclusion criteria

- Adult patients, mentally competent and 18 years of age or older
- Inclusion in the AVARIJN study

Exclusion criteria

- Patients who are not able to give informed consent

Study design

Design

Study type: Observational non invasive

Intervention model: Parallel

Allocation: Non controlled trial

Masking: Open (masking not used)

Control: N/A, unknown

Recruitment

NL

Recruitment status: Pending

Start date (anticipated): 01-07-2016

Enrollment: 59

Type: Anticipated

Ethics review

Not applicable

Application type: Not applicable

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register ID

NTR-new NL5731 NTR-old NTR5918

Other : MEC 16-417

Study results