

KLEM-studie.

No registrations found.

Ethical review	Positive opinion
Status	Recruitment stopped
Health condition type	-
Study type	Observational non invasive

Summary

ID

NL-OMON21641

Source

Nationaal Trial Register

Brief title

KLEM

Health condition

perinatale transmissie van lidocaïne

perinatal transmission of lidocaine

Sponsors and support

Primary sponsor: Amphia Hospital

Source(s) of monetary or material Support: Amphia Hospital

Intervention

Outcome measures

Primary outcome

Plasma concentration lidocaine in the newborn and in the mother on time of delivery.

Secondary outcome

Range of plasma concentration lidocaine in 24 hours in newborn.

Study description

Background summary

Background of the study:

Lidocaine is administered to the mother as local anesthetic before performing a surgically planned incision to the perineum (episiotomy) during delivery.

In Amphibia Hospital two cases have been presented with a possibility of intoxication with lidocaine, following medical signs.

Levels could not exclude an incorrect gift of lidocaine via the maternal perineum directly in the head skin of the child, or a

concentration caused by transmission via the maternal blood when it was administered locally to the mother.

Transmission of lidocaine to the child via the umbilical cord is possible. To define exactly when an intoxication will occur in

newborns, it is necessary to determine the lidocaine concentrations in a child after delivery when lidocaine is locally

administered to the mother. It is also interesting to look at the interval between administration of lidocaine and the time of

delivery in relation to the degree of transmission of lidocaine to the child. In the current literature, no relation could be

confirmed between the concentration lidocaine in the newborn and the interval between administration and partus.

Beside this, it is important to know the pharmacokinetics of lidocaine in newborns. By research the pharmacokinetics of

lidocaine in newborns, we can estimate the duration of exposure to lidocaine and the estimated level of lidocaine

belonging to a specific time.

We aim to investigate the transmission of lidocaine from mother to child during delivery and study the pharmacokinetics of

lidocaine in newborns.

Objective:

With this study we want to determine the concentration lidocaine in newborn after partus using an episiotomy in relation to

the concentration in mother. Our secondary objectives are to look at the relation between the degree of transmission of

lidocaine to the newborns and the drug-delivery interval, and the pharmacokinetical behaviour of lidocaine in newborn

after transmission of lidocaine during the partus.

Study design:

Prospective observational cohort study

Study population:

Fourty obstetrical women with lidocaine for local anesthesia before episiotomy at partus and within this group, all eligible

newborns (minimal 6, maximal 18) who had to stay in the hospital for determine a glucose curve.

Primary study parameters/outcome of the study:

The level lidocaine in the newborn on time of delivery and the level lidocaine in the mother on time of delivery.

Secondary study parameters/outcome of the study:

The range of lidocaine levels in 24 hours in the newborn.

Study objective

There will be a relation between the level lidocaine in newborn, the level in mother and drug-delivery interval.

Study design

1. t=0 hour at delivery;
2. For newborn: t=1, t=3, t=6, t=12, t=24 hour.

Intervention

N/A

Contacts

Public

Amphia Ziekenhuis, Klinische Farmacie

Postbus 90158
J. Huisbrink
Breda 4800 CK
The Netherlands

Scientific

Amphia Ziekenhuis, Klinische Farmacie

Postbus 90158
J. Huisbrink
Breda 4800 CK
The Netherlands

Eligibility criteria

Inclusion criteria

1. Informed consent;
2. Delivery in Amphia Hospital;
3. Administration of lidocaine in the perineum for local anesthesia;
4. Gestation period from 32 weeks;
5. For the newborn: stay in the hospital to determine glucose levels.

Exclusion criteria

1. Administration of lidocaine for other purposes than local anesthesia;
2. Administration of epinefrine together with lidocaine for episiotomy.

Study design

Design

Study type:	Observational non invasive
Intervention model:	Parallel
Allocation:	Non controlled trial
Masking:	Open (masking not used)
Control:	N/A , unknown

Recruitment

NL	
Recruitment status:	Recruitment stopped
Start date (anticipated):	01-01-2013
Enrollment:	40
Type:	Actual

Ethics review

Positive opinion	
Date:	19-12-2012
Application type:	First submission

Study registrations

Followed up by the following (possibly more current) registration

ID: 40012

Bron: ToetsingOnline

Titel:

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register	ID
NTR-new	NL3602
NTR-old	NTR3761
CCMO	NL42283.015.12
ISRCTN	ISRCTN wordt niet meer aangevraagd.
OMON	NL-OMON40012

Study results

Summary results

N/A