

# Cost-effectiveness of IUI, IVF and ICSI for male subfertility. The MAle Subfertility Therapy Effectiveness Rcts.

No registrations found.

|                              |                |
|------------------------------|----------------|
| <b>Ethical review</b>        | Not applicable |
| <b>Status</b>                | Pending        |
| <b>Health condition type</b> | -              |
| <b>Study type</b>            | Interventional |

## Summary

### ID

NL-OMON21655

### Source

Nationaal Trial Register

### Brief title

MASTER study

### Health condition

IVF, IUI, male subfertility (mannelijke subfertiliteit), cost-effectiveness (kosteneffectiviteit)

## Sponsors and support

**Primary sponsor:** Academic Medical Center (AMC)

**Source(s) of monetary or material Support:** ZON-MW, The Netherlands Organization for Health Research and Development

## Intervention

## Outcome measures

### Primary outcome

The primary outcome is establishment of ongoing pregnancy leading to live birth within the treatment time horizon.

## **Secondary outcome**

Time to pregnancy, miscarriage, multiple pregnancy and live birth rate are secondary outcomes. Further secondary outcomes are neonatal mortality, pregnancy complications (preterm birth < 37 weeks, birth weight < 2.500 gram, PIH, (pre-) eclampsia, HELLP) costs of reproductive treatments, perinatal care and ad-verse events. Also patients' quality of life and preferences will serve as secondary outcomes.

# **Study description**

## **Background summary**

Rationale:

We hypothesize that less invasive therapies are equally effective as more invasive therapies for male subfertility.

Objective:

In one third of subfertile couples male subfertility is diagnosed. Current treatments for male subfertility, IUI, IVF and ICSI, have, despite their widespread use, not been compared on their cost-effectiveness. The primary aim of this project is to assess the cost-effectiveness of therapies for male subfertility.

Study design:

IVF versus IUI in moderate male subfertility.

Study population:

Subfertile couples with male subfertility (pre-wash TMSC 3-6 million).

Intervention:

3 cycles of IVF, including transfer of cryoembryos.

Control: 3 cycles of IUI, followed by 3 cycles of IUI-COH. Treatment time horizon 9 months.

Main study parameters/endpoints:

Primary: Ongoing pregnancy leading to live birth.

Secondary: Time to pregnancy, miscarriage, multiple pregnancy, live birth, perinatal outcome, (in-)direct costs, quality of life and patient preferences.

Nature and extent of the burden and risks associated with participation, benefit and group relatedness:

As we compare interventions that are already applied in current practice, no additional risks or burdens are expected from the study.

### **Study objective**

To evaluate the cost-effectiveness of therapies for male subfertility.

### **Study design**

Primary and secondary outcomes within 9 months after randomisation. Questionnaires at one day, two months, four months and six months after randomization.

### **Intervention**

3 cycles of IVF, including transfer of cryopreserved embryos vs 3 cycles of IUI, followed by 3 cycles of IUI-COH.

## **Contacts**

### **Public**

Postbus 90153  
J.P. Bruin, de  
Den Bosch 5200 ME  
The Netherlands  
+31 (0)73 6998660

### **Scientific**

Postbus 90153  
J.P. Bruin, de  
Den Bosch 5200 ME  
The Netherlands  
+31 (0)73 6998660

## Eligibility criteria

### Inclusion criteria

1. Age female partner: 18-38 years;
2. Failure to conceive: 12-36 months;
3. Male subfertility: Pre-wash TMSC  $3-6 \cdot 10^6$ .

### Exclusion criteria

1. Severe male subfertility: Pre-wash TMSC  $< 3 \cdot 10^6$ ;
2. Female partner with polycystic ovary syndrome or any other anovulation, severe endometriosis, double-sided tubal pathology, endocrinopathological disease (Cushing syndrome, adrenal hyperplasia, hyperprolactinemia, acromegaly, imminent ovarian failure, premature ovarian failure, hypothalamic amenorrhea and diabetes mellitus (type I)).

## Study design

### Design

|                     |                             |
|---------------------|-----------------------------|
| Study type:         | Interventional              |
| Intervention model: | Parallel                    |
| Allocation:         | Randomized controlled trial |
| Masking:            | Open (masking not used)     |
| Control:            | Active                      |

## Recruitment

|                           |             |
|---------------------------|-------------|
| NL                        |             |
| Recruitment status:       | Pending     |
| Start date (anticipated): | 01-06-2013  |
| Enrollment:               | 364         |
| Type:                     | Anticipated |

## Ethics review

|                   |                |
|-------------------|----------------|
| Not applicable    |                |
| Application type: | Not applicable |

## Study registrations

### Followed up by the following (possibly more current) registration

No registrations found.

### Other (possibly less up-to-date) registrations in this register

No registrations found.

## In other registers

| Register | ID                                  |
|----------|-------------------------------------|
| NTR-new  | NL3634                              |
| NTR-old  | NTR3822                             |
| Other    | ZonMW : 837002003                   |
| ISRCTN   | ISRCTN wordt niet meer aangevraagd. |

## Study results

### Summary results

N/A