

Subjective well-being, craving for cannabis and compliance or medication switch in a randomized doubleblind study with olanzapine and risperidone.

No registrations found.

Ethical review	Positive opinion
Status	Recruitment stopped
Health condition type	-
Study type	Interventional

Summary

ID

NL-OMON22184

Source

NTR

Brief title

SUB.CAN.OLA.RIS

Intervention

Outcome measures

Primary outcome

1. Subjective Well-being under Neuroleptics scale (SWN);
2. Obsessive Compulsive Drug Use Scale (OCDUS);
3. Positive And Negative Symptoms Scale (PANSS) based on information from the semi-structured interview (SCI- PANSS);
4. Calgary Depression Rating Scale (CDRS);
5. ESRS: Extra pyramidal Symptom Rating Scale;

6. Clinical Global Impression (CGI);
7. Y-BOCS (Yale Brown Obsessive Compulsive Scale);
8. Desires for Drugs Questionnaire (DDQ);
9. Drug Use Self Report (DUSR);
10. Recent Drug Use Urinalysis (RDUU).

Secondary outcome

1. Drop out from the study;
2. Medication compliance and medication switch, symptoms and rehospitalisations during one year follow up, measured with the Life Chart Schedule (LCS).

Study description

Background summary

N/A

Study objective

N/A

Study design

N/A

Intervention

Patients are treated double blind with olanzapine (5-20 mg) or risperidone (1,25-5 mg) for 6 weeks. At t=0, t=7 days and t=42 days, questionnaires are taken and after 6 weeks the medication is disclosed. The physician and patient decide if this neuroleptic will be continued. After one year the questionnaires are taken once more.

Contacts

Public

Erasmus MC
's Gravendijkwal 230
Lonneke Nimwegen, van
Tafelbergweg 25
Rotterdam 3015 CE
The Netherlands
+31 (0)10 4635871

Scientific

Erasmus MC
's Gravendijkwal 230
Lonneke Nimwegen, van
Tafelbergweg 25
Rotterdam 3015 CE
The Netherlands
+31 (0)10 4635871

Eligibility criteria

Inclusion criteria

1. Patients should be able to understand the study description and give informed consent;
2. Diagnosis of schizophrenia, schizoaffective disorder or schizophreniform disorder according to DSM IV;
3. Patients experience a first or second psychotic episode;
4. Age is between 18 and 30 years;
5. No current use of clozapine;
6. Patients must be reliable. They must agree to co operate with all tests and examinations required by the protocol.

Exclusion criteria

1. Pregnancy;
2. Lactating women;
3. Female subject without adequate contraconception;

4. Known hypersensitivity to any ingredient of olanzapine or risperidone;
5. Concomitant use of any other antipsychotic drug than olanzapine or risperidone;
6. Patients are not allowed to have received depot- antipsychotics for a period of at least 3 months prior to the study;
7. Use of other psychotropic medication other than oxazepam or biperiden;
8. Narrow-angle glaucoma;
9. Known neurological or endocrine disease.

Study design

Design

Study type:	Interventional
Intervention model:	Parallel
Allocation:	Randomized controlled trial
Masking:	Double blinded (masking used)
Control:	Active

Recruitment

NL	
Recruitment status:	Recruitment stopped
Start date (anticipated):	01-07-2003
Enrollment:	120
Type:	Actual

Ethics review

Positive opinion	
Date:	22-11-2004
Application type:	First submission

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register	ID
NTR-new	NL4
NTR-old	NTR28
Other	: N/A
ISRCTN	ISRCTN46365995

Study results

Summary results

N/A