Assertive Community Treatment for the elderly.

No registrations found.

Ethical review Positive opinion **Status** Recruiting

Health condition type -

Study type Interventional

Summary

ID

NL-OMON22529

Source

NTR

Brief title

N/A

Health condition

Older psychiatric patients with severe mental disorder who refuse help.

Oudere zorgmijders met ernstige psychiatrische problematiek.

Sponsors and support

Primary sponsor: BavoEuropoort - ParnassiaBavogroep

Source(s) of monetary or material Support: BavoEuropoort - ParnassiaBavogroep

Intervention

Outcome measures

Primary outcome

1. Number of patients who receive care and the number of patients who stay in care;

- 2. The psychiatric-, somatic condition and social functioning of the patient;
- 3. Quality of life.

Secondary outcome

- 1. Ability to live and do things independently;
- 2. Crisis contacts:
- 3. (Court ordered) hospitalization and hospitalization time;
- 4. Increase of motivation for treatment and medication compliance;
- 5. Stronger therapeutic relationship.

Study description

Background summary

A randomised controlled trial examines the effect of Assertive Community Treatment for older psychiatric patients with seriously various problems who refuse help compared with care as usual.

Study objective

As compared to care as usual Assertive Community Treatment for the elderly creates:

- 1. Dropout prevention;
- 2. Decrease in psychiatric symptoms;
- 3. Decrease in somatic symptoms;
- 4. Decrease in social problems;
- 5. Increase in Quality of Live.

Study design

Baseline, 9 months after randomisation and 18 months after randomisation.

Intervention

Assertive Community Treatment.

Contacts

Public

Erasmus MC, Universitair Medical Centre Rotterdam, department psychiatry, DP-0122 PObox 2040

J. Stobbe Rotterdam 3000 CA The Netherlands +31(0)10 7032355

Scientific

Erasmus MC, Universitair Medical Centre Rotterdam, department psychiatry, DP-0122 PObox 2040

J. Stobbe Rotterdam 3000 CA The Netherlands +31(0)10 7032355

Eligibility criteria

Inclusion criteria

- 1. 65 years or older;
- 2. patients with a psychiatric diagnosis (or strong supposed diagnosis) on Axis I of the DSM IV-TR;
- 3. lack of motivation for treatment;
- 4. regular treatment is not sufficient;
- 5. seriously various problems.

Exclusion criteria

- 1. Moderate to seriously dementia;
- 2. Simple addiction.

Study design

Design

Study type: Interventional

Intervention model: Parallel

Allocation: Randomized controlled trial

Masking: Single blinded (masking used)

Control: Active

Recruitment

NL

Recruitment status: Recruiting
Start date (anticipated): 01-03-2008

Enrollment: 160

Type: Anticipated

Ethics review

Positive opinion

Date: 08-01-2009

Application type: First submission

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register ID

NTR-new NL1549 NTR-old NTR1620 Other : N/A

ISRCTN wordt niet meer aangevraagd

Study results

Summary results

N/A