Alveolar ridge preservation with a xenograft and a collagen matrix or a free connective tissue graft versus spontaneous healing: A 1-year prospective randomized clinical trial.

No registrations found.

Ethical review Positive opinion **Status** Recruiting

Health condition type -

Study type Interventional

Summary

ID

NL-OMON22864

Source

Nationaal Trial Register

Brief title

Mucograft Seal Study

Health condition

Oral implants, alveolar ridge preservation

Sponsors and support

Catharina Hospital Eindhoven, Department Oral and Maxillofacial Surgery

Source(s) of monetary or material Support: Geistlich Pharma AG, Wolhusen,

Switzerland

Straumann AG, Basel, Switzerland

Intervention

Outcome measures

Primary outcome

The level of the buccal marginal gingiva (one year after implant loading)

Secondary outcome

- 1. Peri-implant esthetic score (PES)
- 2. White esthetic score (WES)
- 3. Labial soft tissue volume (using impressions)
- 4. The marginal bone level around the implants/ Distance from implant shoulder to first bone-to-implant contact (MBL/DIB) using a standardized digital intra-oral radiograph
- 5. Evaluation of facial bone wall (using CBCT)
- 6. The plaque index (PI)
- 7. The bleeding index (B)
- 8. The gingiva index (GI)
- 9. The pocket probing depth (PPD)
- 10. The width of the attached mucosa (WAM)
- 11. Patient's satisfaction about the esthetical result of the crown and peri-implant tissue

Tertiary: Implant succes

Study description

Background summary

Replacement of a single tooth in the esthetic zone is a demanding procedure. The application of a biomaterial in the extraction socket, covered with a collagen matrix or a soft tissue graft

2 - Alveolar ridge preservation with a xenograft and a collagen matrix or a free con ... 24-05-2025

may lead to less vertical and horizontal changes of the alveolar ridge and soft-tissues and thereby simplifying the procedure with more predictable outcomes.

The study is a prospective randomized clinical trial with 1-year follow-up. Patients in need for an implant-supported dental crown to replace a maxillary tooth in the esthetic zone are randomized in one of three techniques: A) bone substitute material and a collagen matrix B) bone substitute material covered with a palatal graft, or C) spontaneous healing. After extraction of the tooth, patients will be treated according to their assigned protocol. After 8 weeks a Straumann implant will be placed. The implants are loaded after a minimum healing time of approximately 8-10 weeks. Esthetic and clinical parameters and patient satisfaction is assessed after tooth extraction, before implant placement and up to one year after crown placement. Esthetic scores consist of the Peri-implant esthetic score (PES) and the White esthetic score (WES). Labial soft tissue volume is assessed using digitized casts; the buccal bone is assessed using Cone Beam CT scans. Other assessments are the buccal marginal gingiva, marginal bone level (MBL), plague index (PI), the bleeding index BI, the gingiva index (GI), the pocket probing depth (PPD) and the width of the attached mucosa (WAM). A patient's questionnaire includes a visual analog scales (VAS) that will focus on expectation and satisfaction of the surgical procedure and about the esthetic result of the dental crown and the peri-implant tissue.

Study objective

Alveolar ridge preservation with a xenograftand a collagen matrix or a free connective tissue graft, compared to spontaneous healing, leads to better soft tissue and bone volume for early implant placement and a better aesthetic result, up to 1 year after functional loading.

Study design

Pre-operative, one week post extraction, implant placement, 2 weeks after implant placement 1, 6 and > 12 months after placement of the crown.

Intervention

Group A) Placement of a bone substitute material (deproteinized bovine bone mineral with 10% collagen; Bio-Oss Collagen®, Geistlich) covered with a collagen matrix (Geistlich Mucograft® Seal).

Group B) Placement of a bone substitute material (deproteinized bovine bone mineral with 10% collagen; Bio-Oss Collagen®, Geistlich) covered with a punch biopsy of the palate.

Group C) Spontaneous healing (control group).

Contacts

Public

Department of Oral & Maxillofacial Surgery, Special Dental Care and Orthodontics

B.P. Jonker
Erasmus MC Rotterdam
P.O. Box 2040, 3000 CA Rotterdam, the Netherlands
's-Gravendijkwal 230, Office D-224
Rotterdam
The Netherlands
Tel + 31 10 703 4138

Scientific

Department of Oral & Maxillofacial Surgery, Special Dental Care and Orthodontics

B.P. Jonker
Erasmus MC Rotterdam
P.O. Box 2040, 3000 CA Rotterdam, the Netherlands
's-Gravendijkwal 230, Office D-224
Rotterdam
The Netherlands
Tel + 31 10 703 4138

Eligibility criteria

Inclusion criteria

Over 18 years of age. Need for an implant-supported dental crown to replace a maxillary tooth at the location of an incisor, cuspid or first/second bicuspid; single tooth diastema as a maximum; intact buccal bone plate (confirmed by clinical examination); sufficient occlusal and mesio-distal dimensions for insertion of one implant with a functional prosthetic restoration.

Exclusion criteria

Presence of clinical active periodontal disease; presence of an acute inflammatory oral disease; smoking; uncontrolled diabetes; a history of radiotherapy in the head- and-neck region or current chemotherapy; disability (mental and/or physical) to maintain basic oral hygiene procedures; under eighteen years of age.

Study design

Design

Study type: Interventional

Intervention model: Parallel

Allocation: Randomized controlled trial

Masking: Single blinded (masking used)

Control: Active

Recruitment

NL

Recruitment status: Recruiting
Start date (anticipated): 09-11-2015

Enrollment: 75

Type: Anticipated

Ethics review

Positive opinion

Date: 09-09-2017

Application type: First submission

Study registrations

Followed up by the following (possibly more current) registration

ID: 42186

Bron: ToetsingOnline

Titel:

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register ID

NTR-new NL6497 NTR-old NTR6685

CCMO NL49965.078.14 OMON NL-OMON42186

Study results

Summary results

N/A