Physiotherapy, is there a difference? Consequence of less movement in patients with ankylosing spondylitis.

No registrations found.

Ethical review Positive opinion

Status Recruitment stopped

Health condition type -

Study type Observational non invasive

Summary

ID

NL-OMON23026

Source

Nationaal Trial Register

Brief title

FSA

Health condition

ankylosing spondylitis, physiotherapy, mobility

Sponsors and support

Primary sponsor: Atrium Medical Center and Zuyd Hogeschool Heerlen

Source(s) of monetary or material Support: Initiator

Intervention

Outcome measures

Primary outcome

Consequence on the mobility, quality of live, disease activity and VAS in patients with

1 - Physiotherapy, is there a difference? Consequence of less movement in patients ... 24-05-2025

akylosing spondylitis due to not paying for the consult of the physiotherapist by the health insurance of the patient.

Methods of measurement: Questionnaires VAS, SF36, BASDAI, BASFI, movement diary and use of painmedication.

Secondary outcome

Use of pain medication.

Study description

Background summary

Physiotherapy is there a difference?

Consequence of less movement in patients with ankylosing spondylitis.

Consequence on the mobility with patients who have akylosing spondylitis due to not paying for the consult of the physiotherapist by the health insurance of the patient.

The patient will fill in serveral questionnaires at 3 or 4 moments prospective in time.

Study objective

Consequence on the mobility of patients who have akylosing spondylitis due to not paying for the consult of the physiotherapist by the health insurance of the patient.

Study design

Start aug/ september 2012 until april/ may 2013.

Intervention

N/A

Contacts

Public

Henri Dunantstraat 5 K. Pintaric Heerlen 6419 PC The Netherlands +31 (0)45 5766688

Scientific

Henri Dunantstraat 5 K. Pintaric Heerlen 6419 PC The Netherlands +31 (0)45 5766688

Eligibility criteria

Inclusion criteria

- 1. Patient with ankylosing spondylitis;
- 2. Patient 18 years or older;
- 3. Patient who have had physiotherapy;
- 4. Patient willing to give informed consent.

Exclusion criteria

Patients with a recent excacerbation of ankylosing spondylitis.

Study design

Design

Study type: Observational non invasive

Intervention model: Parallel

Allocation: Non-randomized controlled trial

Masking: Open (masking not used)

Control: Active

Recruitment

NL

Recruitment status: Recruitment stopped

Start date (anticipated): 01-08-2012

Enrollment: 75

Type: Actual

Ethics review

Positive opinion

Date: 24-07-2012

Application type: First submission

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register ID

NTR-new NL3401 NTR-old NTR3544 Other : 12-N-75/III

ISRCTN wordt niet meer aangevraagd.

Study results

Summary results

N/A