

# Treatment outcomes for gestational diabetes diagnosed either according to WHO 2013 or WHO 1999 thresholds: the TANGO-DM Randomized Controlled Trial

Published: 06-09-2018

Last updated: 14-09-2024

We hypothesized that intensive GDM treatment according to the new (2013) GDM criteria would result in a reduction in infants with birth weight >90th centile (large for gestational age, LGA), in comparison to treatment according to the old...

<b>Ethical review</b>	Approved WMO
<b>Status</b>	Recruitment stopped
<b>Health condition type</b>	Maternal complications of pregnancy
<b>Study type</b>	Interventional

## Summary

### ID

NL-OMON23396

### Source

NTR

### Brief title

TANGO-DM

### Condition

- Maternal complications of pregnancy

### Health condition

GDM Diabetes gravidarum Gestational diabetes zwangerschapsdiabetes

### Research involving

Fetus in utero, Human

## Sponsors and support

**Primary sponsor:** Leading the Change

**Source(s) of monetary or material Support:** Leading the Change

## Intervention

- Food (substances)

## Explanation

## Outcome measures

### Primary outcome

Large for gestational age (LGA, defined as birth weight >90th centile, using the Dutch Perined reference charts)

### Secondary outcome

Secondary perinatal outcomes will include:

- Perinatal mortality (up until 6 weeks postpartum)
- Birth weight
- Birth weight >4000 g
- Small for gestational age ( • Gestational age at delivery
- Preterm delivery (<37 weeks)
- Reason for preterm delivery (iatrogenic, spontaneous)
- 1-min Apgar score
- 5-min Apgar score
- 10-min Apgar score
- Shoulder dystocia
- Birth injury (i.e. fractures, nerve palsy)
- Cord blood C peptide (in a subset, where available)
- Neonatal glucose postpartum: 1-3-6-12-24 hours or similar (in GDM treatment group) • Neonatal hypoglycaemia (moderate, serum glucose <2.6 mmol/l; severe, serum glucose <2.0 mmol/l)
- Neonatal intensive care/medium care admission
- Hyperbilirubinemia necessitating phototherapy
- Respiratory support > 24 hours
- Neonatal encephalopathy

Secondary outcomes of delivery:

- Mode of delivery (spontaneous, assisted vaginal, caesarean). We will split instrumental deliveries for fetal distress or non-progressive labour
- Induction of labour
- Post-partum haemorrhage

Secondary maternal outcomes will include:

- Hypertensive disorders of pregnancy (including preeclampsia and pregnancy induced hypertension)
- Need for treatment with insulin or oral glucose lowering medication
- Maternal total weight gain in gestation
- Glycemic control: diary (in GDM treatment group)
- Puerperal sepsis requiring antibiotics
- Maternal admission postpartum
- Serious health outcomes up to the time of primary hospital discharge assessed from patient medical records<sup>23</sup>.
- Intention to breastfeed prior to giving birth
- Breastfeeding at the end of postpartum period

Cost effectiveness analysis:

- Societal costs
- Quality-adjusted life years

Both from a societal and healthcare perspective

## Study description

### Background summary

Gestational diabetes mellitus (GDM), or hyperglycemia first diagnosed in pregnancy, affects 7-10% of all pregnancies worldwide. Perinatal risk rises with increasing glycemia at oral glucose tolerance test (OGTT). The new (2013) WHO criteria recommend a lower fasting, and a higher post-load threshold for GDM diagnosis in comparison to the old (1999) WHO criteria. To date, however, outcomes of GDM treatment for those affected by the altered diagnostic criteria, has not been well investigated.

### Study objective

We hypothesized that intensive GDM treatment according to the new (2013) GDM criteria would result in a reduction in infants with birth weight >90th centile (large for gestational age, LGA), in comparison to treatment according to the old criteria (1999).

### Study design

Multicentre open-label randomized controlled trial (RCT).

### Intervention

The intervention under investigation will be intensive GDM treatment of women with OGTT results discordant between the new and the old criteria. Treatment will comply with the NVOG guideline: monitoring of blood glucose, dietary recommendations and pharmacotherapy for those unable to achieve euglycemia with dietary intervention alone.

The comparator will be routine obstetric care.

## Contacts

**Public**

**Scientific**

## Eligibility criteria

### Age

Adults (18-64 years)

Adults (18-64 years)

Elderly (65 years and older)

Elderly (65 years and older)

### Inclusion criteria

- Singleton pregnancy
- Aged >18 years
- OGTT for all indications
- OGTT conducted between gestational ages 16+0 and 30+0 weeks
- Discordant result on a 3-point 75-gram OGTT i.e.:
  - o Fasting glucose > 5.1 and <7.0 mmol/l OR
  - o 1-hour glucose ≥10.0 mmol/l OR
  - o 2-hour glucose >7.8 and <8.4 mmol/l
- Gestational age <32+0 at study inclusion
- Written informed consent

### Exclusion criteria

- known preconception diabetes
- major fetal congenital /chromosomal abnormality (eg trisomy 21, spina bifida), known at time of randomization
- significant medical or psychiatric co-morbidities as judged by the investigator (e.g. high dose corticosteroid treatment)
- inability to understand written informed consent without help as indicated by their usual care provider

## Study design

### Design

Study phase:	N/A
Study type:	Interventional
Intervention model:	Parallel
Allocation:	Randomized controlled trial
Masking:	Open (masking not used)
Control:	N/A , unknown
Primary purpose:	Diagnostic

### Recruitment

NL	
Recruitment status:	Recruitment stopped
Start date (anticipated):	04-10-2018
Enrollment:	1032
Type:	Actual

### IPD sharing statement

**Plan to share IPD:** Undecided

## Ethics review

Approved WMO	
Date:	16-08-2018
Application type:	First submission
Review commission:	MEC Academisch Medisch Centrum (Amsterdam)
	Kamer G4-214
	Postbus 22660
	1100 DD Amsterdam
	020 566 7389

## Study registrations

### Followed up by the following (possibly more current) registration

ID: 52951

Bron: ToetsingOnline

Titel:

### Other (possibly less up-to-date) registrations in this register

No registrations found.

### In other registers

Register	ID
NTR-new	NL7127
NTR-old	NTR7473
CCMO	NL63013.018.18
OMON	NL-OMON52951

## Study results