# Heart Failure and Promotion of Physical Activity before and after Cardiac reHabilitation

No registrations found.

Ethical review	Positive opinion
Status	Pending
Health condition type	-
Study type	Interventional

# **Summary**

### ID

NL-OMON24242

**Source** Nationaal Trial Register

**Brief title** HF-aPProACH

#### **Health condition**

Heart Failure, Patients with chronic HFrEF (ischaemic or non-ischaemic)

### **Sponsors and support**

**Primary sponsor:** Erasmus Medical Center **Source(s) of monetary or material Support:** H2020-Bigmedilytics

#### Intervention

### **Outcome measures**

#### **Primary outcome**

6MWT

#### Secondary outcome

1 - Heart Failure and Promotion of Physical Activity before and after Cardiac reHabi ... 29-05-2025

• Maintained PA (step count) after CR, • Difference in enrollment percentage in CR, • Hospitalization and mortality, • Health related quality of life, • Physical fitness:, • Questionnaires: Mental health (HADS), Fear of movement (TSK-NL Heart), • Compliance to program

# **Study description**

### **Background summary**

Rationale: Heart failure is a severe chronic disease with a great health care burden. Frequent (re-)hospitalizations and a high mortality remain an important issue. Optimal medical therapy (OMT) and lifestyle changes such as increased physical activity (PA) are the cornerstones of treatment. Based on the European Society for Cardiology (ESC) guidelines, regular aerobic exercise, for example within a centre-based cardiac rehabilitation (CR) is recommended for all Heart Failure patients with reduced Ejection Fraction (HFrEF) patients. Several studies have shown that PA is just as effective as medical therapy and can lower hospital admissions and decrease mortality. However, physical activity in this group of patients is challenging as both low participation rates for CR and relapse to low physical activity levels after CR are major issues. This study aims to demonstrate that monitoring physical activity in combination with motivational feedback benefits the level of participation in centre-based CR and, hence, the outcomes for these HF patients.

Objective: To investigate in our defined HF patient population if monitoring in combination with motivation on physical activity leads to an increase in enrollment rate for cardiac rehabilitation and sustained physical activity after cardiac rehabilitation as measured by the 6MWT.

Study design: The proposed study is a randomized controlled trial with a follow-up of at least 6 months. A total of 180 patients will be randomized to 2 arms in a 2:1 fashion: (1) Physical activity monitoring device with feedback and motivation; (2) Physical activity monitoring device without feedback and motivation. Both arms will have standard of care (SoC), including standard cardiac rehabilitation (specific for HF) according to the Dutch guidelines along with OMT as prescribed by the treating physician. The enrolment period will be 12 months. Additional measurements will take place at 3 time points: at baseline (within 6 weeks after inclusion), at the end of CR (16-20 weeks after inclusion) and at the end of follow-up (3 months after CR or at least 6 months after inclusion).

Study population: Patients aged 18-85 years with chronic HFrEF (NYHA class II and III) who have a clinically stable condition, an indication for physical exercise and CR, will be eligible to participate in this study. All included patients are required to have provided written informed consent.

Intervention: All patients in both arms of the study will wear an Actigraph activity tracker for physical activity monitoring. In addition, only the intervention arm will also wear a Fitbit smartwatch on which they will receive automated motivational feedback based on a goal for physical activity (step count), set by the treating physician before and after CR.

### **Study objective**

2 - Heart Failure and Promotion of Physical Activity before and after Cardiac reHabi ... 29-05-2025

We hypothesize that physical activity (PA) monitoring with motivational feedback (using an activity tracker) before and after centre-based CR in patients with HFrEF (NYHA class II/III) will lead to a clinically meaningful increase in the 6MWT. Contributing factors to this objective are an expected improved participation rate in CR and a maintained PA after CR.

#### Study design

• T0: At inclusion.

• T1: At completion\* of CR, or for patients who did not start or did not complete CR, this will be at 16-20 weeks after inclusion.

• T2: 3 months after completion of CR, or for patients who did not start or did not complete CR, this will be 7 months after inclusion.

#### Intervention

The intervention arm will be stimulated to become more active. This will be achieved through visualization of step count on the Fitbit smartwatch and mobile phone and by receiving feedback on daily step count goals on the Fitbit smartwatch and mobile phone.

# Contacts

**Public** Erasmus MC Arne IJpma

0107044656 Scientific Erasmus MC Arne IJpma

0107044656

# **Eligibility criteria**

## **Inclusion criteria**

1. Age  $\geq$  18 years,  $\leq$  85 years.

2. Patients with chronic HFrEF (ischaemic or non-ischaemic) from the outpatient clinic.

3. NYHA class II/III.

4. Indication for and referral to CR (first time or last CR was more than 1 year before inclusion).

3 - Heart Failure and Promotion of Physical Activity before and after Cardiac reHabi ... 29-05-2025

- 5. Clinically stable (no hospitalization in the month prior to inclusion).
- 6. At least 1 cardiology-related hospitalization in 1 -12 months prior to inclusion.

7. The patient should be willing and able to cooperate with study procedure (use activity tracker/device, fill out questionnaires, comply with the required follow-up visits and sign informed consent.

8. Sufficient understanding of Dutch language, both spoken and written.

9. Access to a smartphone with SMS functionality and a personal computer with internetaccess and an email address.

### **Exclusion criteria**

- 1. Patients with acute HF, HFpEF or HFmEF.
- 2. Patients who are currently or have been (< 1 month) hospitalized.
- 3. End stage renal disease requiring dialysis.
- 4. Other comorbidities/conditions precluding exercise training.
- 5. Patients with a life expectancy <1 year.
- 6. Patients on the waiting list for a heart transplant or other major surgery planned <1 year.
- 7. Patients already actively using their own smart watch.

# Study design

## Design

Study type:	Interventional
Intervention model:	Parallel
Allocation:	Randomized controlled trial
Masking:	Open (masking not used)
Control:	Active

# Recruitment

NL	
Recruitment status:	Pending
Start date (anticipated):	02-12-2019
Enrollment:	180
Туре:	Anticipated

### **IPD** sharing statement

Plan to share IPD: No

# **Ethics review**

Positive opinionDate:28-11-2019Application type:First submission

# **Study registrations**

# Followed up by the following (possibly more current) registration

ID: 49839 Bron: ToetsingOnline Titel:

### Other (possibly less up-to-date) registrations in this register

No registrations found.

### In other registers

Register	ID
NTR-new	NL8190
ССМО	NL68986.078.19
OMON	NL-OMON49839

# **Study results**