

# The role of Ketotofen in the treatment of postoperative ileus.

No registrations found.

<b>Ethical review</b>	Positive opinion
<b>Status</b>	Recruitment stopped
<b>Health condition type</b>	-
<b>Study type</b>	Interventional

## Summary

### ID

NL-OMON24494

### Source

Nationaal Trial Register

### Brief title

N/A

### Intervention

### Outcome measures

#### Primary outcome

Gastric emptying.

#### Secondary outcome

Symptoms.

## Study description

### Background summary

This is a randomized, placebo-controlled, double-blind study with patients who'll undergo

abdominal surgery. They will be treated with Ketotofen 4 mg or 12 mg or placebo per day. The patient will start his medication intake 3 days before surgery till 2 days after surgery. Post operative effectivity will be determined by scintigraphy to look at gastric retention and colontransit. This scintigraphy will take place 24 hours after surgery. The patients will be clinical evaluated by filling out a daily symptom score list.

### **Study objective**

N/A

### **Study design**

N/A

### **Intervention**

Ketotofen 4 mg or 12 mg or placebo per day.

## **Contacts**

### **Public**

Academic Medical Center (AMC), Department of Gastroenterology, C2-328,  
P.O. Box 22660  
G.E.E. Boeckxstaens  
Meibergdreef 9  
Amsterdam 1100 DD  
The Netherlands  
+31 (0)20 5667375

### **Scientific**

Academic Medical Center (AMC), Department of Gastroenterology, C2-328,  
P.O. Box 22660  
G.E.E. Boeckxstaens  
Meibergdreef 9  
Amsterdam 1100 DD  
The Netherlands  
+31 (0)20 5667375

## **Eligibility criteria**

## Inclusion criteria

Laparotomy for (malignant) proces originaty from colon or female reproduction organs.

## Exclusion criteria

Pre-operative radiation, intake of medication with effect on the Gastrointestinal motility, intake of immunosuppressiva, intake of anti-allergy medication, intra-abdominal inflammation (cholecystitis or absces included).

## Study design

### Design

Study type:	Interventional
Intervention model:	Parallel
Allocation:	Randomized controlled trial
Masking:	Double blinded (masking used)
Control:	Placebo

### Recruitment

NL	
Recruitment status:	Recruitment stopped
Start date (anticipated):	01-04-2004
Enrollment:	48
Type:	Actual

## Ethics review

Positive opinion	
Date:	02-06-2005
Application type:	First submission

## Study registrations

### Followed up by the following (possibly more current) registration

No registrations found.

### Other (possibly less up-to-date) registrations in this register

No registrations found.

### In other registers

Register	ID
NTR-new	NL16
NTR-old	NTR38
Other	: N/A
ISRCTN	ISRCTN34927143

## Study results

### Summary results

N/A