

A study on the effect of information on the use of analgesics in patients after weight loss surgery.

No registrations found.

Ethical review	Positive opinion
Status	Pending
Health condition type	-
Study type	Interventional

Summary

ID

NL-OMON25083

Source

NTR

Brief title

NUBS

Health condition

bariatric surgery; use of NSAIDS
bariatrische chirurgie; gebruik van NSAID's

Sponsors and support

Primary sponsor: Medisch Centrum Leeuwarden (MCL), Leeuwarden, The Netherlands
Medisch Centrum Leeuwarden (MCL), H. Dunantweg 2, 8934 AD Leeuwarden

Source(s) of monetary or material Support: Wetenschapsfonds MCL, Leeuwarden, The Netherlands

Intervention

Outcome measures

Primary outcome

The percentage of users of NSAIDS in the period 6 months before the intervention and in the period 3 to 9 months after the intervention in comparison with the control group. The use is determined by collecting dispensing data from pharmacies 6 months before the intervention and 9 months after the intervention. The use of NSAIDS is to be considered as stopped if there is no use of an NSAID any more in the period 3 to 9 months after the intervention.

Secondary outcome

The number of Defined Daily Doses (DDDS) of an NSAID in the period 6 months before the intervention and in the period 3 to 9 months after the intervention in comparison to the control group. The number of DDDS is determined by means of the amount of NSAIDS dispensed in the period 6 months before and 3 to 9 months after the intervention. The intervention is to be considered effective, if the mean number of DDDS of an NSAID in the period 3 to 9 months after the intervention is reduced by half in comparison with the mean number of DDDS of an NSAID in the period 6 months before the intervention.

The number of users of NSAIDS with no concurrent use of a PPI in the period 6 months before the intervention and in the period 3 to 9 months after the intervention in comparison to the control group.

There is concurrent use of a PPI with an NSAID if the period of use of a PPI has a $\geq 2/3$ overlap with the period of use of an NSAID. The use is determined by collecting dispensing data from pharmacies 6 months before and 9 months after the intervention. The intervention is to be considered effective if the number of users of NSAIDS with no concurrent use of a PPI is reduced by half in the period 3 to 9 months after the intervention in comparison with the period 6 months before the intervention.

Study description

Background summary

Background of the study:

Although U.S. guidelines indicate that the use of NSAIDS should be avoided in patients after bariatric surgery, a significant portion of the population of patients who underwent bariatric surgery in Medical Center Leeuwarden (MCL) in the period October 2008 to December 2011, use an NSAID at some time after the bariatric surgery. Moreover in the MCL only half of the patients use a PPI while using an NSAID after bariatric surgery.

These data indicate that there is room for improvement in the prescribing and use of NSAIDS in patients who have undergone bariatric surgery.

Objective of the study:

Primary objective:

To determine the effect of an intervention on the level of reduction of the number of users of NSAIDS in patients who underwent bariatric surgery in Medisch Centrum Leeuwarden (MCL).

Secondary objectives:

To determine the effect of an intervention on the level of reduction of the use of NSAIDS in patients who underwent bariatric surgery in MCL. To study the use of proton pump inhibitors (PPIS) in patients who use an NSAID after bariatric surgery.

Study design:

A randomized controlled intervention study in patients who underwent bariatric surgery in MCL between March 2011 and November 2011. There will be a nine months follow-up period after intervention.

Study population:

Patients who underwent bariatric surgery in MCL between March 2011 and November 2011.

Intervention:

The intervention is performed by sending a letter to patients and their family doctors belonging to the intervention group, with information about the risks of use of NSAIDS by patients after bariatric surgery. As a control, a group of patients is included in which no intervention is performed.

Primary study parameters/outcome of the study:

The percentage of users of NSAIDS in the period 6 months before the intervention and in the period 3 to 9 months after the intervention in comparison with the control group. The use is determined by collecting dispensing data from pharmacies 6 months before the intervention and 9 months after the intervention. The use of NSAIDS is to be considered as stopped if there is no use of an NSAID any more in the period 3 to 9 months after the intervention.

Secondary study parameters/outcome of the study:

The number of Defined Daily Doses (DDDS) of an NSAID in the period 6 months before the intervention and in the period 3 to 9 months after the intervention in comparison to the control group. The number of DDDS is determined by means of the amount of NSAIDs dispensed in the period 6 months before and 3 to 9 months after the intervention. The intervention is to be considered effective, if the mean number of DDDS of an NSAID in the period 3 to 9 months after the intervention is reduced by half in comparison with the mean number of DDDS of an NSAID in the period 6 months before the intervention.

The number of users of NSAIDs with no concurrent use of a PPI in the period 6 months before the intervention and in the period 3 to 9 months after the intervention in comparison to the control group.

There is concurrent use of a PPI with an NSAID if the period of use of a PPI has a $\geq 2/3$ overlap with the period of use of an NSAID. The use is determined by collecting dispensing data from pharmacies 6 months before and 9 months after the intervention.

Nature and extent of the burden and risks associated with participation, benefit and group relatedness:

Safety issues are not applicable for this study. The current policy of the MCL is that prior to surgery bariatric surgeons inform patients that NSAIDs after bariatric surgery should be avoided whenever possible. Patients in the control group only undergo a randomization and then the regular procedure. For those patients this is only an observational study. Patients in the intervention group will get more information about the risks associated with the use of NSAIDs after bariatric surgery. The subsequent part of the study in patients in the intervention group, is also only observational. The burden and risks for patients participating in this study, in the intervention group, as well as the control group are very low. Two times during the study dispensing data are requested from pharmacies and nine months after the intervention all patients will get a telephone call in which a questionnaire is used to record OTC use of NSAIDs and indications for NSAID use.

Study objective

Although U.S. guidelines indicate that the use of NSAIDs should be avoided in patients after bariatric surgery, a significant portion of the population of patients who underwent bariatric surgery in Medical Center Leeuwarden (MCL) in the period October 2008 to December 2011, use an NSAID at some time after the bariatric surgery. These data indicate that there is room for improvement in the prescribing and use of NSAIDs and PPIs in patients who have undergone bariatric surgery. Informing patients and their family doctors about the risks of use of NSAIDs after bariatric surgery should reduce the use of NSAIDs.

Primary objective of the study:

To determine the effect of an intervention on the level of reduction of the number of users of NSAIDS in patients who underwent bariatric surgery in Medisch Centrum Leeuwarden (MCL).

Secondary objectives:

To determine the effect of an intervention on the level of reduction of the use of NSAIDS in patients who underwent bariatric surgery in MCL.

To determine the use of protonpump inhibitors in patients who use NSAIDS after bariatric surgery.

The intervention is performed by sending a letter to patients and their family doctors belonging to the intervention group, with information about the risks of use of NSAIDS by patients after bariatric surgery. As a control, a group of patients is included in which no intervention is performed.

Study design

1. Before intervention;
2. Intervention;
3. 9 months after intervention.

Intervention

The intervention is performed by sending a letter to patients and their family doctors belonging to the intervention group, with information about the risks of use of NSAIDS by patients after bariatric surgery.

As a control, a group of patients is included in which no intervention is performed.

Two times during the study dispensing data are requested from pharmacies and nine months after the intervention all patients will get a telephone call in which a questionnaire is used to record OTC use of NSAIDS and indications for NSAID use.

Contacts

Public

Medisch Centrum Leeuwarden

Henri Dunantweg 2
J.P. Yska
Leeuwarden 8934 AD
The Netherlands

+31 (0)58 2866604

Scientific

Medisch Centrum Leeuwarden

Henri Dunantweg 2

J.P. Yska

Leeuwarden 8934 AD

The Netherlands

+31 (0)58 2866604

Eligibility criteria

Inclusion criteria

1. Patients who underwent bariatric surgery in Medisch Centrum Leeuwarden (MCL) in the period March to November 2011;
2. Patients who signed and turned in a written informed consent.

Exclusion criteria

Patients with no medication record at any moment of the study.

Study design

Design

Study type:	Interventional
Intervention model:	Parallel
Allocation:	Randomized controlled trial
Masking:	Single blinded (masking used)
Control:	N/A , unknown

Recruitment

NL	
Recruitment status:	Pending
Start date (anticipated):	01-11-2012

Enrollment: 240
Type: Anticipated

Ethics review

Positive opinion
Date: 15-10-2012
Application type: First submission

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register	ID
NTR-new	NL3485
NTR-old	NTR3665
Other	: RTPO 877
ISRCTN	ISRCTN wordt niet meer aangevraagd.

Study results

Summary results

N/A