

Determining the tumor uptake of labelled bevacizumab in children with high grade or diffuse intrinsic pontine glioma on PET scans.

No registrations found.

Ethical review	Positive opinion
Status	Recruiting
Health condition type	-
Study type	Observational non invasive

Summary

ID

NL-OMON27519

Source

NTR

Brief title

89Zr-Bmab PET in pHGG & DIPG

Health condition

Diffuse intrinsic pontine glioma

Pediatric high grade glioma

Malignant glioma

Hooggradig glioom

Ponglioom

Sponsors and support

Primary sponsor: VU University Medical Center

Intervention

Outcome measures

Primary outcome

VEGF-expression measured by Standard Uptake Values of ⁸⁹Zr-Bevacizumab in pHGG and DIPG.

Secondary outcome

1. Optimal moment of scanning obtained by five patients with positive ⁸⁹Zr-bevacizumab uptake;
2. Body biodistribution and dosimetry of ⁸⁹Zr-bevacizumab.

Study description

Background summary

Paediatric high grade gliomas (pHGG) including diffuse intrinsic pontine gliomas (DIPG) have a poor prognosis. PET imaging with labelled antibodies enables drug distribution investigations and non-invasive target expression studies. pHGG and DIPG highly express vascular endothelial growth factor (VEGF) on RNA level, which is involved in mitogenic, angiogenic, and permeability enhancing processes. Monoclonal antibody bevacizumab inhibits VEGF-A and showed efficacy in adult glioma and to a lesser extend in pHGG. Bevacizumab is labelled to Zirconium-89, a positron emitter with a long half-time which is preferable because of its safety, purity and stable binding to its antibody and relatively low costs. In adults, ⁸⁹Zr-bevacizumab could be used safely in humans and was shown to visualise targets precisely. In this study, bevacizumab is administered in a microdose at 1/100th of the therapeutic dose in pHGG and DIPG. PET scans are performed at 1, 72 and 144 hours post-injection. We expect that PET-imaging of ⁸⁹Zr-bevacizumab may help to select patients more likely to respond to bevacizumab therapy.

Study objective

Is there VEGF-expression in DIPG & HGG measured by the tumor uptake of Zr-bevacizumab?

Study design

⁸⁹Zr-bevacizumab is injected and PET scans are performed at 1, 72 and 144 hours post-injection.

Intervention

This is a diagnostic PET study. The labelled antibody is ⁸⁹Zr-Bevacizumab.

Contacts

Public

VU University Medical Center, Department of Pediatric Oncology/Hematology,
De Boelelaan 1117
Gertjan J.L. Kaspers
De Boelelaan 1117
Amsterdam 1081 HV
The Netherlands
+31 (0)20 4442420

Scientific

VU University Medical Center, Department of Pediatric Oncology/Hematology,
De Boelelaan 1117
Gertjan J.L. Kaspers
De Boelelaan 1117
Amsterdam 1081 HV
The Netherlands
+31 (0)20 4442420

Eligibility criteria

Inclusion criteria

1. DIPG (MRI confirmed, biopsy not required) de novo;
2. De novo biopsy proven HGG patients with minimal residual tumor of 0.5 mm in each dimension or;
3. pHGG & DIPG patients with progressive disease after radiotherapy;
4. Age between 4 and 18 years;
5. Able to lay down quiet for 30 minutes.

Exclusion criteria

1. Chemotherapy or radiotherapy in the past two weeks;

2. Previous administration of bevacizumab or another anti-VEGF drug;
3. Known hypersensitivity against humanized monoclonal antibodies;
4. Neurofibromatosis type I.

Study design

Design

Study type:	Observational non invasive
Intervention model:	Parallel
Allocation:	Non controlled trial
Masking:	Open (masking not used)
Control:	N/A , unknown

Recruitment

NL	
Recruitment status:	Recruiting
Start date (anticipated):	01-09-2011
Enrollment:	15
Type:	Anticipated

Ethics review

Positive opinion	
Date:	09-07-2012
Application type:	First submission

Study registrations

Followed up by the following (possibly more current) registration

ID: 38105
Bron: ToetsingOnline

Titel:

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register	ID
NTR-new	NL3370
NTR-old	NTR3518
CCMO	NL34922.000.11
ISRCTN	ISRCTN wordt niet meer aangevraagd.
OMON	NL-OMON38105

Study results

Summary results

Jansen MH, van Vuurden DG, Vandertop WP, Kaspers GJ. Diffuse intrinsic pontine gliomas: A systematic update on clinical trials and biology. Cancer Treat Rev. 2012;38:27-35

Van Dongen GA, Visser GW, Lub-de Hooge MN, de Vries EG, Perk LR. Immuno-PET: a navigator in monoclonal antibody development and applications. Oncologist 2007;12:1379-1389.