Well-being therapy in outpatients with mild affective disorders

No registrations found.

Ethical review Positive opinion

Status Pending

Health condition type -

Study type Interventional

Summary

ID

NL-OMON28550

Source

NTR

Health condition

Well-being therapy Anxiety disorders Depression Short-term treatment Positive psychology

Welbevinden therapie Angststoornis Depressie Kortdurende behandeling Positieve psychologie

Sponsors and support

Primary sponsor: Dimence, a mental health care centre. The study is being performed in the course of the education of gz-psychologist to clinical psychologist.

Source(s) of monetary or material Support: Dimence, a mental health care centre. The study is being performed in the course of the education of gz-psychologist to clinical psychologist.

Intervention

Outcome measures

Primary outcome

Well-being, as measured with the Mental Health Continuum-Short Form (MHC-SF). Psychological complaints as measured with the Hospital Anxiety and Depression Scale (HADS).

Secondary outcome

The experience of patients and therapists with well-being therapy.

Study description

Study objective

Firstly this study tries to answer the question whether applying well-being therapy to outpatients with a mild affective disorder in Dutch mental health care, reveals a trend in which well-being is increased and psychiatric complaints are reduced.

Secondly the intention is to learn more about the experiences of the participants (patients as well as therapists) with this new form of therapy, to be able to carry out further development and adjustments if necessary.

The outcome of this study can be used to account for a randomized controlled trail.

Study design

Participants will be treated with well-being therapy. They will be asked to answer two questionnaires at three specific moments (T0, T1, T2). In total participants will spend a maximum of 75 minutes on the questionnaires over a period of 6 months. Also, one in three participants will randomly be invited to participate in a semi-structured interview.

Intervention

Participants will be treated with well-being therapy, a short-term protocol (6 sessions) in which they will actively focus on the enhancement of well-being.

Contacts

Public

Ropcke Zweerslaan 2

E.J.B. Peeneman Unknown The Netherlands 0523-280280

Scientific

Ropcke Zweerslaan 2

E.J.B. Peeneman Unknown The Netherlands 0523-280280

Eligibility criteria

Inclusion criteria

- -Classification according to DSM IV-TR of an anxiety or affective disorder as follows from an intake.
- -Participants are indicated for the 'middle' category in mental health care.
- -A below average score (<2.13) on the Mental Health Continuüm-Short Form (MHC- SF) before treatment. (this cut off score was based on the mean minus one SD (Lamers et al., 2011) and the expectation that clients with a higher score will have less possibilities for measurable improvements.
- -Age between 18 and 65 years.

Exclusion criteria

- Poor Dutch language.
- Complex psychiatric problems which requires special mental health care.

Study design

Design

Study type: Interventional

Intervention model: Parallel

Allocation: Randomized controlled trial

Control: N/A, unknown

Recruitment

NL

Recruitment status: Pending

Start date (anticipated): 01-05-2015

Enrollment: 33

Type: Anticipated

Ethics review

Positive opinion

Date: 01-04-2015

Application type: First submission

Study registrations

Followed up by the following (possibly more current) registration

ID: 42118

Bron: ToetsingOnline

Titel:

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register ID

NTR-new NL4983 NTR-old NTR5128

CCMO NL50778.044.15 OMON NL-OMON42118

Study results