

Nierdenervatie als behandeling van hoge bloeddruk na niertransplantatie.

No registrations found.

Ethical review	Not applicable
Status	Suspended
Health condition type	-
Study type	Interventional

Summary

ID

NL-OMON29017

Source

NTR

Brief title

CRESCENT

Health condition

Therapy resistant hypertension
renal allograft recipients

Sponsors and support

Primary sponsor: Department of Nephrology
Academic Medical Center at the University of Amsterdam

Source(s) of monetary or material Support: Nederlandse Nierstichting
other applications pending

Intervention

Outcome measures

Primary outcome

Change daytime blood pressure after 6 months after renal denervation assessed by ambulatory blood pressure measurement; compared to continued standard care (i.e.

conforming to National Kidney Foundation Kidney Disease Quality Outcomes Quality Initiative guidelines).

Secondary outcome

1. Change in 123I-MIBG uptake of native kidneys (i.e. effectiveness of denervation);
2. Change in muscle sympathetic nerve activity, plasma catecholamines, rennin and aldosterone;
3. Change in proteinuria after 6 months;
4. Change in eGFR after 6 months;
5. Change in number of antihypertensive drugs;
6. Change in health related quality of life.

Study description

Background summary

Rationale:

In patients with a renal allograft, hypertension is a major etiological factor for cardiovascular morbidity, mortality and allograft nephropathy. Controlling hypertension in patients with a renal allograft is therefore crucial. There is a pressing, yet currently unmet clinical need for new blood pressure lowering strategies in renal allograft recipients. The diseased native kidneys are major contributors to hypertension, through neuro-hormonal up-regulation that leads to high levels of renin and sympathetic activity. Recently a catheter-based approach has been developed to disrupt renal sympathetic nerves. Currently this innovative technique has only been tested to lower blood pressure in therapy resistant hypertensive patients without significant renal disease. We hypothesize that catheter based selective renal denervation of the native kidneys in renal transplant recipients will improve blood pressure control and diminish the number of antihypertensive drugs.

Objective:

To test the efficacy and safety of renal sympathetic denervation therapy with a special focus on preservation of renal allograft function.

Study design:

We propose a randomized controlled clinical trial (intervention group n=20; controls n=20). Intervention and control groups will receive standard protocolized antihypertensive treatment prior to, and during the trial. The intervention group will receive renal denervation in addition to standard treatment.

Study population:

Inclusion criteria are a renal allograft in situ since > 6 months with an estimated GFR >35 ml/min/ 1,73m² and a diuresis > 200ml/day of the native kidneys at time of transplantation (to ensure presence of vital neuro-hormonally active kidney tissue) and a daytime blood pressure >140/90 mmHg (assessed by 24-hours ambulatory measurement) while on >3 antihypertensive medications in maximal tolerated dose, including a diuretic.

Intervention:

Prior to study-inclusion all patients will receive standard protocolized hypertension treatment based on the National Kidney Foundation Kidney Disease Quality Outcomes Quality Initiative guidelines (2004). Renal sympathetic denervation is achieved by the interventional radiologist percutaneously entering the lumen of the main renal artery of each of the native kidneys, with a catheter connected to a radiofrequency generator. He applies 6-8 radiofrequency ablations within each renal artery. The procedure is performed in an outpatient clinic setting and patients receive standard measures for prevention of contrast nephropathy.

Main study parameters/endpoints:

Primary endpoint is blood pressure reduction after 6 months (day time blood pressure assessed by 24-hours ambulatory measurement). Secondary outcomes include changes in renal sympathetic innervation (by ¹²³I-MIBG scintigraphy), systemic sympathetho-humoral activity (by peroneal microneurography and plasma catecholamines and rennin and aldosterone activity), eGFR, proteinuria, number of anti-hypertensive drugs needed and quality of life and adverse events.

Study objective

We hypothesize that catheter based selective renal denervation of the native kidneys in renal transplant recipients will improve blood pressure control and diminish the number of antihypertensive drugs.

Study design

Primary and secondary measurement: At inclusion and after 6 months.

Intervention

Catheter based renal denervation of the native kidneys (Symplicity system, Medtronic). Renal nerve ablation is achieved in a single 40 minute catheterisation session.

The control group will receive care as usual.

Contacts

Public

Department of Nephrology

Academic Medical Center at the University of Amsterdam
C.T.P. Krediet
Amsterdam
The Netherlands

Scientific

Department of Nephrology

Academic Medical Center at the University of Amsterdam
C.T.P. Krediet
Amsterdam
The Netherlands

Eligibility criteria

Inclusion criteria

1. Renal graft in situ since > 6 months, estimated GFR >35 ml/min/1,73m² and;
2. Diuresis of the native kidneys at transplant >200 ml/day (to ensure the presence of vital kidney that could be affected by renal nerve ablation) and;
3. Day time blood pressure >140/90 mmHg (assessed by 24-hours ambulatory measurement within 3 month prior to inclusion in the study, as is regularly performed in the nephrology outpatient clinic) while;
4. Treated according to National Kidney Foundation Kidney Disease Quality Outcomes Quality Initiative guidelines (2004), i.e. having been advised to minimize salt intake and using >3

antihypertensive medications in maximal tolerated dose, including a diuretic. Medications and their dosages should not have been changed since the measurement.

Exclusion criteria

1. (Planned) pregnancy, lactation;
2. Life expectancy < 1 year;
3. Contraindications for (relative) hypotensive episodes i.e. haemodynamically significant valvular disease, documented transient ischaemic attacks or angina pectoris during relative hypotension;
4. Complications during previous radiological interventions (i.e. allergy to contrast agent, cholesterol embolism);
5. (Reno)vascular abnormalities in any part of the catheter access route (including severe femoral or renal artery stenosis and atherosclerosis, previous renal stenting or angioplasty, or known dual renal arteries);
6. Use of vitamine K antagonists or other (non-aspirin) form of anti-coagulatory therapy.

Study design

Design

Study type:	Interventional
Intervention model:	Parallel
Allocation:	Randomized controlled trial
Masking:	Open (masking not used)
Control:	N/A , unknown

Recruitment

NL	
Recruitment status:	Suspended
Start date (anticipated):	01-11-2011
Enrollment:	40
Type:	Anticipated

Ethics review

Not applicable

Application type:

Not applicable

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register	ID
NTR-new	NL2856
NTR-old	NTR2998
Other	ABR : 37711
ISRCTN	ISRCTN wordt niet meer aangevraagd.

Study results

Summary results

N/A