Efficiency of the acupuncture in the treatment of the pain of neck of origin myofascial.

No registrations found.

Ethical review Not applicable

Status Recruitment stopped

Health condition type -

Study type Interventional

Summary

ID

NL-OMON29504

Source

Nationaal Trial Register

Brief title

N/A

Health condition

Persons with pain of neck and head of muscle origin.

Persons who works as administrative officers and use the computer frequently.

Persons with not traumatic pain of more of three months of evolution

Personen mit Schmerz des Halses und Hauptes von Muskelursprung.

Personen, der als administrative Offiziere und Gebrauch der Computer häufig arbeitet.

Personen mit nicht traumatischer Schmerz von mehr von drei Monaten der Entwicklung

Sponsors and support

Primary sponsor: It does not exist

Source(s) of monetary or material Support: We haven't funding

Intervention

Outcome measures

Primary outcome

After four weeks.

Secondary outcome

After three mounths.

Study description

Background summary

N/A

Study objective

The treatment of the muscle lower trapezius improves the pain of affected persons of problems craneocervicales of origin miofascial.

Study design

- 1. Four meetings, once per week;
- 2. One assement after four weeks;
- 3. Second assement after three mounths.

Intervention

- 1. The first, assement the cervical spine mobility, muscle pain and algometry of lower trapezius PGM;
- 2. The second, dry needle of PGM of lower trapezius;
- 3. The third, assement the cervical spine mobility, muscle pain and algometry of lower trapezius PGM, again.

Contacts

Public

c/Camilo José Cela, 53 - Portal A - 1ºC 28806 - Alcalá de Henares Madrid

Physiotherapy Departament External Campus, s/n Daniel Pecos Martín Physiotherapy Departament External Campus, s/n Madrid 28871 E The Netherlands

Scientific

c/Camilo José Cela, 53 - Portal A - 1ºC 28806 - Alcalá de Henares Madrid

Physiotherapy Departament External Campus, s/n Daniel Pecos Martín Physiotherapy Departament External Campus, s/n Madrid 28871 E The Netherlands

Eligibility criteria

Inclusion criteria

- 1. 18-year-old major being;
- 2. Pain presents cranium-cervical of three months of evolution;
- 3. The criteria fulfill diagnosis described for the triggers points.

Exclusion criteria

- 1. Existence of some contraindication for the invasive treatment:
- A. Ingestion of some anticoagulant;
 - 3 Efficiency of the acupuncture in the treatment of the pain of neck of origin myo ... 14-05-2025

- B. Allergy to the nickel;
- C. To have suffered some neumotorax;
- D. Fear of the needles.
- 2. To present diagnosis of fibromialgia or to have widespread pain;
- 3. To present diagnosis of migraine of type tensional or migraine;
- 4. To have suffered a cervical traumatism;
- 5. To be diagnosed of an injury of cervical sprain;
- 6. To present some inflammatory artropatía;
- 7. To be controlled of cancer of breast;
- 8. To present a linfedema;
- 9. To be pregnant;
- 10. To have been an agreement with acupuncture in the last six months;
- 11. To have received invasive treatment for the PGM in the last six months.

Study design

Design

Study type: Interventional

Intervention model: Parallel

Allocation: Randomized controlled trial

Masking: Double blinded (masking used)

Control: Placebo

Recruitment

NL

Recruitment status: Recruitment stopped

Start date (anticipated): 03-10-2008

4 - Efficiency of the acupuncture in the treatment of the pain of neck of origin myo ... 14-05-2025

Enrollment: 90

Type: Actual

Ethics review

Not applicable

Application type: Not applicable

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register ID

NTR-new NL1873 NTR-old NTR1987

Other :

ISRCTN wordt niet meer aangevraagd.

Study results

Summary results

N/A