

Endoscopic Tri-Modal Imaging (ETMI) for the detection of early neoplasia in patients with Barrett's esophagus (BE) in tertiary referral centers; a randomized cross-over multi-center study.

Published: 23-01-2007

Last updated: 14-05-2024

Does ETMI improve the dysplasia in Barrett's esophagus?

Ethical review	-
Status	Recruitment stopped
Health condition type	Malignant and unspecified neoplasms gastrointestinal NEC
Study type	Observational invasive

Summary

ID

NL-OMON30430

Source

ToetsingOnline

Brief title

ETMIcross

Condition

- Malignant and unspecified neoplasms gastrointestinal NEC

Synonym

Barrett's esophagus, esophageal cancer

Research involving

Human

Sponsors and support

Primary sponsor: Academisch Medisch Centrum

Source(s) of monetary or material Support: Ministerie van OC&W,Olympus, Japan,Olympus;Japan

Intervention

Keyword: Autofluorescence, Barrett's esophagus, Early neoplasia, Narrow Band Imaging

Outcome measures

Primary outcome

Number of patients detected with dysplasia.

Number of lesions detected with dysplasia.

Secondary outcome

Positive predictive value of high-resolution white light endoscopy and autofluorescence imaging.

Reduction of fals-positive findings with narrow band imaging.

Study description

Background summary

Barrett's esophagus is a premalignant disease. Dysplastic change occur in a sequential order from low grade dysplasia - high grade dysplasia - intramucosal carcinoma and finally to advanced esophageal adenocarcinoma. Patients with Barrett's esophagus undergo regular endoscopic surveillance for the detection of dysplasia. Dysplastic lesions are, however, difficult to detect with standard endoscopy techniques.

A new prototype, Endoscopic TriModal Imaging, incorporates high-resolution white light endoscopy, autofluorescence imaging and narrow band imaging in one system. Autofluorescence imaging excites the esophagus with blue light. The mucosa fluorescence and normal tissue has a different fluorescent spectrum than dysplastic tissue. In uncontrolled studies it was found that autofluorescence may improve the detection of dysplasia. This technique was however also associated with false-positive findings. Narrow band imaging is a novel technique that makes use of special light filters that improve the imaging of superficial mucosal structures and blood vessels in the mucosa. Regular mucosal and vascular patterns are associated with non-dysplastic Barrett's and irregular patterns with dysplastic Barrett's. After the detection of suspicious areas with

autofluorescence narrow band imaging can be used for the detailed inspection of these areas and thus subsequently lower the number of false positive findings. Patients with an early detected dysplastic/neoplastic lesion(s) can be treated curatively endoscopically, making an esophagectomy obsolete.

Study objective

Does ETMI improve the dysplasia in Barrett's esophagus?

Study design

A randomised cross-over multi-center study. Patients will undergo two endoscopies, one with ETMI and with the standard technique. Both procedures will be conducted by two different endoscopists.

Study burden and risks

An endoscopic procedure is safe. Most complications are related with the sedation during the procedure. Total morbidity of a gastroscopy is between 0.14% and 0.20%. Mortality is 0.07%. The additional risk of a second procedure is very small.

Contacts

Public

Academisch Medisch Centrum

Meibergdreef 9

1105 AZ

Nederland

Scientific

Academisch Medisch Centrum

Meibergdreef 9

1105 AZ

Nederland

Trial sites

Listed location countries

Netherlands

Eligibility criteria

Age

Adults (18-64 years)

Elderly (65 years and older)

Inclusion criteria

Age >18 years

Prior diagnosis of Barrett's esophagus

Minimum Barrett's length of C>2M>2 or C<2M>4 according to the Prague C&M classification

Written informed consent

Exclusion criteria

Presence of esophagitis > grade A according to the LA classification of erosive esophagitis

Presence of conditions precluding histological sampling of the esophagus

Study design

Design

Study type:	Observational invasive
Intervention model:	Parallel
Allocation:	Randomized controlled trial
Masking:	Open (masking not used)
Control:	Active
Primary purpose:	Diagnostic

Recruitment

NL

Recruitment status: Recruitment stopped

Start date (anticipated):	01-01-2006
Enrollment:	31
Type:	Actual

Ethics review

Not available

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register	ID
CCMO	NL15477.018.06