

# The effect of post-thoracotomy pulmonary rehabilitation on quality of life.

Published: 22-03-2007

Last updated: 10-05-2024

Primary: To evaluate the effects of a rehabilitation program on health related quality of life (SGRQ) in the 12 months postoperative period in patients with an elective thoracotomy. Secondary: To evaluate the effects of a rehabilitation program on...

<b>Ethical review</b>	Approved WMO
<b>Status</b>	Recruiting
<b>Health condition type</b>	Other condition
<b>Study type</b>	Interventional

## Summary

### ID

NL-OMON30490

### Source

ToetsingOnline

### Brief title

Post-thoracotomy rehabilitation and quality of life.

### Condition

- Other condition
- Respiratory tract neoplasms
- Economic and housing issues

### Synonym

Post-thoracotomy rehabilitation; lung-exercise program after thoraxsurgery.

### Health condition

longrevalidatie

### Research involving

Human

## Sponsors and support

**Primary sponsor:** Isala Klinieken

**Source(s) of monetary or material Support:** maatschap ongziekten en isala klinieken.

## Intervention

**Keyword:** pain, quality of life, rehabilitation, thoracotomy

## Outcome measures

### Primary outcome

1. health related quality of life (SGRQ score).

### Secondary outcome

1. general quality of life (SF-36).
2. acute / chronic post-thoracotomy pain (McGill and VAS).
3. impairment (changes in pulmonary function).
4. disability (exercise capacity; 6 Minute Walking Distance).
5. start to complete recovery (ECOG score of 0 or 1).

## Study description

### Background summary

Morbidity in the post operative phase of thoracotomies is characterised by pain, dyspnea, shoulder dysfunction and a loss of exercise tolerance may occur. There is literature about post-thoracotomy painsyndromes and postoperative quality of life has been investigated. However, data on post-thoracotomy rehabilitation and influence on morbidity and recovery are not available.

### Study objective

Primary:

To evaluate the effects of a rehabilitation program on health related quality

of life (SGRQ) in the 12 months postoperative period in patients with an elective thoracotomy.

Secondary:

To evaluate the effects of a rehabilitation program on general quality of life (SF-36), acute / chronic postthoracotomy pain, impairment (changes in pulmonary function), disability (exercise capacity) and start to complete recovery (ECOG score of 0 or 1) 12 months postoperatively in patients with an elective thoracotomy.

## **Study design**

Eighty-eight elective thoracotomy patients with an age between the 18 and 80 years will be recruited during two years. The follow up for each patient will be one year. The exclusion criteria are chronic pain involvement, a previous thoracotomy, comorbidity limiting rehabilitation, psychiatric illness and non-compliance.

Patients will be asked for participation before thoracotomy by informed consent and will be randomised post-thoracotomy, before discharge into a rehabilitation or 'regular care' group. Regular care is the regular approach after discharge; without a pulmonary rehabilitation program and with permission of regular physical therapy or when indicated (not related to thoracotomy). Regular sporting and exercises (no special training programs) will be tolerated. Randomisation will take place for rehabilitation or regular care by a minimisation program with special attendance to age ( $< 70$  or  $\geq 70$  years), gender, the result of the 6 minute walk test ( $< 100$  m or  $\geq 100$  m), the FEV1 ( $FEV1 < 40\%$  or  $\geq FEV1 40\%$ ) and type of surgery (pneumectomy vs other). During screening a spirometry and a 6 minute walk test will be performed. The first questionnaires (VAS, McGill pain questionnaire) will be given to the patients after enrolment and before thoracotomy. The SGRQ and the SF 36 will be given before, during and after rehabilitation.

The post-thoracotomy pain management consists of the standard analgetic treatment; a thoracic epidural catheter which will be slowly replaced by paracetamol, opioids and NSAID's. The consumption of analgetics will also be documented in the follow up. Supplementary oxygen and lung inhalation is given when necessary. This will be scored too. Regular physical therapy will be started directly after thoracotomy as soon as possible.

The rehabilitation program will be initiated within a month after discharge. The degree of the rehabilitation program will be separately detected for each patient by a cycle test (heart rate) within 3 weeks post-thoracotomy. Patients in the pulmonary rehabilitation program will be guided by a multidisciplinary team of pulmonologist, physical therapist and social worker during 12 weeks. The program will last 3 x 2 hours weekly consisting of exercise training and

education.

All patients will be followed up after discharge at 1 month, 3 months, 6 months and 12 months at the outpatient clinic of the pulmonology department and the pain clinic of the anaesthesia department. Before discharge and at these intervals (except for the SGRQ at 1 month), patients have to fill questionnaires about quality of life and pain (SF 36, SGRQ; McGill pain questionnaire and VAS). Exercise capacity by means of the 6 minutes walk test and spirometry will be measured after the accomplishment of the rehabilitation program at 3 months. Patients of the control group will also be asked during this visits whether they have had regular physical therapy or whether they are sporting to evaluate the exact effects of the rehabilitation program.

During follow up all the results between the experimental rehabilitation group and the regular care group will be compared. In the interpretation of these results start to complete recovery is defined as an ECOG score of 0 or 1.

## **Intervention**

An early pulmonary rehabilitation program during 12 weeks (3x2 hours) initiated within a month after discharge from the hospital and will be guided by a multidisciplinary team consisting of pulmonologist, social worker and physical therapist.

## **Study burden and risks**

Burden and risks associated with participation:

There are no adverse effects due to the rehabilitation program. Patients in the \*regular care\* group will not be guided by the multidisciplinary rehabilitation team.

The disadvantages are the following: Patients have to fill in quality of life questionnaires (SGRQ and SF-36), have to fill in pain questionnaires, have to evaluate a painscale at the pain clinic (VAS and McGill Pain questionnaire) and have to perform the 6 minutes walk-test and spirometry tests. Finally, patients have to visit the pulmonologist and the pain clinic more frequently.

The expected benefits are a faster recovery with a better exercise tolerance, less pain and a better quality of life.

## Contacts

### Public

Isala Klinieken

Groot Wezenland 20  
8011 J W Zwolle  
Nederland

### Scientific

Isala Klinieken

Groot Wezenland 20  
8011 J W Zwolle  
Nederland

## Trial sites

### Listed location countries

Netherlands

## Eligibility criteria

### Age

Adults (18-64 years)

Elderly (65 years and older)

### Inclusion criteria

1. Elective thoracotomy patients.
2. Age  $\geq 18 \leq 80$  years.
3. ECOG 0-2 post-thoracotomy.

### Exclusion criteria

1. Patients with chronic pain.
2. A previous thoracotomy.
3. Comorbidity limiting rehabilitation:
  - RA
  - severe ischaemic heart disease or myocardial failure; EF  $\leq 35$  %.

- muscle diseases
  - fibromyalgia
  - neurological disorders (Parkinson, CVA and laesions of the spinal cord).
  - psychiatric diseases
4. Non-compliance.

## Study design

### Design

Study type:	Interventional
Intervention model:	Parallel
Allocation:	Randomized controlled trial
Masking:	Open (masking not used)

**Primary purpose:** Other

### Recruitment

NL	
Recruitment status:	Recruiting
Start date (anticipated):	01-04-2007
Enrollment:	88
Type:	Actual

## Ethics review

Approved WMO	
Date:	22-03-2007
Application type:	First submission
Review commission:	METC Isala Klinieken (Zwolle)

## Study registrations

**Followed up by the following (possibly more current) registration**

No registrations found.

## Other (possibly less up-to-date) registrations in this register

No registrations found.

## In other registers

Register	ID
CCMO	NL14089.075.06