

Activating the frontal brain in schizophrenia: comparison of aripiprazole versus risperidone using functional magnetic resonance imaging

Published: 26-06-2007

Last updated: 08-05-2024

This study looks if the third generation antipsychotic aripiprazole can improve activity of the prefrontal cortex and cognitive and social function, when compared to the second generation antipsychotic risperidone.

Ethical review	Approved WMO
Status	Recruitment stopped
Health condition type	Schizophrenia and other psychotic disorders
Study type	Observational invasive

Summary

ID

NL-OMON31229

Source

ToetsingOnline

Brief title

MRI study on the effect of Abilify versus Risperdal

Condition

- Schizophrenia and other psychotic disorders

Synonym

chronic psychosis, Schizophrenia

Research involving

Human

Sponsors and support

Primary sponsor: Universitair Medisch Centrum Groningen

Source(s) of monetary or material Support: European Young Investigators Award

Intervention

Keyword: aripiprazole, fMRI, frontal cortex, risperidone

Outcome measures

Primary outcome

Difference in brain activity of the prefrontal cortex between groups with different medication. Besides accuracy and reaction times during task performance and severity of positive and negative symptoms.

Secondary outcome

nvt

Study description

Background summary

Schizophrenia is characterized by a widespread cognitive dysfunction, affecting the domains of memory, learning, attention and executive functioning. Besides, disabilities in social functioning exist, one of the main characteristics of schizophrenia. These problems are possibly caused by diminished activity of the prefrontal cortex. The third generation antipsychotic Abilify (aripiprazole) can restore/improve the activity of the prefrontal cortex. This could cause a decrease in negative symptoms and improve social cognition. Risperdal (risperidone) has another mode of action, in which dopamine is blocked in the whole brain. Therefore the expectations are that activity of the prefrontal cortex will not increase substantially when Risperdal is prescribed, in contrast to Abilify that will activate the prefrontal cortex.

Study objective

This study looks if the third generation antipsychotic aripiprazole can improve activity of the prefrontal cortex and cognitive and social function, when compared to the second generation antipsychotic risperidone.

Study design

24 patients suffering from schizophrenia will be randomly assigned to Abilify (aripiprazole) or Risperdal (risperidone), only when no clear preference exists for one of the two kinds of medication. During the treatment there will be regular visits to the clinician in charge. Before onset of medication intake an MRI experiment will be conducted. Subsequently a transition period of three weeks to Abilify or Risperdal takes place. In this period the current medication is gradually decreased while the intake of the new medicines is gradually increased. Next treatment with new medication will be continued for four weeks. After this period a second MRI experiment will take place. The fMRI experiments are composed of different tasks which activate the prefrontal cortex. Activity of the prefrontal cortex will be compared between patient groups, before and after use of Abilify and Risperdal. Activity of the prefrontal cortex and social cognition should improve (more) after treatment with aripiprazole. Besides positive and negative symptoms of schizophrenia will be rated with a structured interview, the PANSS. Besides, 20 healthy controls will be included. They will be subjected to one MRI-session, and no antipsychotics will be administered. On a separate occasion they will be interviewed with the mini-SCAN. This is a short, diagnostic interview to exclude psychiatric illnesses.

Study burden and risks

The study consists of two MRI experiments with a scantime of 75 minutes, a structured interview (PANSS, 30 minutes) and two questionnaires. Comparable studies have been carried out before and were experienced as only being a slight effort for subjects.

The medication that is used is prescribed by a clinician only if this is medically relevant. In case of side effects or increase of symptoms the study will be terminated.

Contacts

Public

Universitair Medisch Centrum Groningen

Antonius Deusinglaan 2
9713 AW Groningen
NL

Scientific

Universitair Medisch Centrum Groningen

Antonius Deusinglaan 2
9713 AW Groningen

Trial sites

Listed location countries

Netherlands

Eligibility criteria

Age

Adults (18-64 years)

Elderly (65 years and older)

Inclusion criteria

People diagnosed for schizophrenia according to DSM-IV. They should not respond to first choice treatment or suffer from serious side effects.

Besides, healthy control without high education level.

Exclusion criteria

Subjects with a psychiatric or neurologic disease other than schizophrenia, for which they have been treated. Presence of MRI incompatible implants. For safety reasons female participants who may be pregnant will be excluded.

Study design

Design

Study type:	Observational invasive
Intervention model:	Parallel
Allocation:	Randomized controlled trial
Masking:	Open (masking not used)
Control:	Active

Primary purpose: Other

Recruitment

NL

Recruitment status: Recruitment stopped

Start date (anticipated): 10-01-2008

Enrollment: 44

Type: Actual

Medical products/devices used

Product type: Medicine

Brand name: Abilify

Generic name: Aripiprazole

Registration: Yes - NL intended use

Product type: Medicine

Brand name: Risperdal

Generic name: Risperidone

Registration: Yes - NL intended use

Ethics review

Approved WMO

Date: 26-06-2007

Application type: First submission

Review commission: METC Universitair Medisch Centrum Groningen (Groningen)

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register	ID
EudraCT	EUCTR2007-002748-79-NL
CCMO	NL17987.042.07