

# The prevalence of postpartum posttraumatic stress symptoms, depression, anxiety and the need of control in women delivered in The Netherlands

Published: 20-09-2007

Last updated: 08-05-2024

To assess the prevalence of postpartum posttraumatic stress symptoms, depression, anxiety and the need of control in women delivered in The Netherlands.

|                              |   |
|------------------------------|---|
| <b>Ethical review</b>        | Approved WMO  |
| <b>Status</b>                | Pending   |
| <b>Health condition type</b> | Pregnancy, labour, delivery and postpartum conditions |
| <b>Study type</b>            | Observational non invasive                            |

## Summary

### ID

NL-OMON31262

### Source

ToetsingOnline

### Brief title

Postpartum psychiatric disorders

### Condition

- Pregnancy, labour, delivery and postpartum conditions

### Synonym

Psychiatric disorders postpartum

### Research involving

Human

## Sponsors and support

**Primary sponsor:** Academisch Medisch Centrum

**Source(s) of monetary or material Support:** Ministerie van OC&W

## Intervention

**Keyword:** anxiety, delivery, depression, post traumatic stress

## Outcome measures

### Primary outcome

Post partum stress symptoms

### Secondary outcome

anxiety, depression, personality, sense of coherence, the need for control and the extend of control

## Study description

### Background summary

Traumatic reactions to childbirth are an important public health issue. Risk factors have been identified. Many of them could be detected during pregnancy. However, a suitable instrument is lacking. Also it is not clear how to intervene to reduce the traumatic reactions to childbirth. Anticipating on a prospective study on this issue, in this retrospective cohort study the prevalence will be measured of postpartum posttraumatic stress symptoms, depression, anxiety and the need of control in women delivered in The Netherlands 1. at home, 2. in a rural hospital and 3. in a university hospital

### Study objective

To assess the prevalence of postpartum posttraumatic stress symptoms, depression, anxiety and the need of control in women delivered in The Netherlands.

### Study design

Retrospective cohort design

## Study burden and risks

It takes about 60 minutes to fill in the questionnaires on the website or in print

## Contacts

### Public

Academisch Medisch Centrum

Hanzeplein 1  
9700 RB Groningen  
Nederland

### Scientific

Academisch Medisch Centrum

Hanzeplein 1  
9700 RB Groningen  
Nederland

## Trial sites

### Listed location countries

Netherlands

## Eligibility criteria

### Age

Adults (18-64 years)  
Elderly (65 years and older)

### Inclusion criteria

Three months postpartum

### Exclusion criteria

Not being able to speak Dutch

## Study design

### Design

**Study type:** Observational non invasive

Masking: Open (masking not used)

Control: Uncontrolled

Primary purpose: Prevention

### Recruitment

NL

Recruitment status: Pending

Start date (anticipated): 01-05-2007

Enrollment: 700

Type: Anticipated

## Ethics review

Approved WMO

Application type: First submission

Review commission: METC Universitair Medisch Centrum Groningen (Groningen)

## Study registrations

### Followed up by the following (possibly more current) registration

No registrations found.

### Other (possibly less up-to-date) registrations in this register

No registrations found.

**In other registers**

| Register | ID             |
|----------|----------------|
| CCMO     | NL16448.042.07 |