

Perioperative temperature management: a big small problem

Published: 08-02-2008

Last updated: 10-05-2024

Does preoperative heating of patients undergoing elective orthopedic surgery reduce the incidence of intraoperative hypothermia and postoperative comfort and modulate postoperative glucose and insulin blood levels?

Ethical review	Approved WMO
Status	Recruitment stopped
Health condition type	Other condition
Study type	Observational invasive

Summary

ID

NL-OMON31705

Source

ToetsingOnline

Brief title

HOT (HypOThermia)

Condition

- Other condition
- Bone and joint therapeutic procedures

Synonym

Perioperative hypothermia, perioperative temperature loss

Health condition

Perioperatief bloedverlies en postoperatieve hyperglycemie

Research involving

Human

Sponsors and support

Primary sponsor: Vrije Universiteit Medisch Centrum

Source(s) of monetary or material Support: Ministerie van OC&W

Intervention

Keyword: Hypothermia, Prewarming, Recovery, Surgery

Outcome measures

Primary outcome

Incidence of perioperative hypothermia

Secondary outcome

Perioperative blood loss

Patient discomfort as defined by pain, nausea and vomiting, shivering and thermal discomfort

Postoperative glucose and insulin levels

Study description

Background summary

Hypothermia, defined as a core temperature below 36 degrees Celcius, is commonly seen in patients undergoing surgery. Intraoperative hypothermia is related to postoperative patient discomfort, impaired wound healing, increased blood loss, anxiety and disturbances in glucose and insulin levels. Prewarming of patients prior to surgery may result in maintenance of intraoperative temperature and reduce postoperative complications. We therefore aim to investigate the effects of prewarming of patients undergoing elective orthopedic surgery on intraoperative temperature control and blood loss and postoperative patient discomfort and glucose and insulin levels.

Study objective

Does preoperative heating of patients undergoing elective orthopedic surgery reduce the incidence of intraoperative hypothermia and postoperative comfort

and modulate postoperative glucose and insulin blood levels?

Study design

Multi-center, prospective, randomized clinical trial.

Study burden and risks

All measurement parameters are part of standard perioperative and clinical care, except for drawing of blood samples for the determination of glucose, insulin, Hb, Ht, creatinine kinase, lactate, APTT and INR (in total 76 ml)

There are three modalities that will be applied to draw patient blood:

1) Blood sampling from an intravenous catheter. Peripheral intravenous catheter placement is standard perioperative procedure in all surgical patients, and will therefore not add up to patient discomfort in the present study.

2) Blood sampling by intravenous puncture. There is a 10-20% chance that blood sampling can not sufficiently be performed through an intravenous catheter. In this case, blood sampling will be performed by an intravenous puncture. Blood sampling will be performed on three different time-points, two during anesthesia and one at the postoperative ward. On the postoperative ward patients are recovering from anesthesia and patient discomfort due to blood sampling will be the highest in this particular population.

3) Glucose levels using a glucose meter device. Postoperative serum glucose levels can be measured using a glucose meter device. A single puncture in the index finger to obtain blood will be necessary and this will be minimal invasive.

Contacts

Public

Vrije Universiteit Medisch Centrum

De Boelelaan 1117

1081 HV

Nederland

Scientific

Vrije Universiteit Medisch Centrum

De Boelelaan 1117

1081 HV

Nederland

Trial sites

Listed location countries

Netherlands

Eligibility criteria

Age

Adults (18-64 years)

Elderly (65 years and older)

Inclusion criteria

Patients undergoing an elective orthopedic knee- or hip replacement

Participation based on informed consent

> 18 years old and < 85 years old

Exclusion criteria

BMI $18.5 < x < 40$

Pregnancy

Non elective trauma patients

Infectious or febrile patients ($> 37.5^{\circ}\text{C}$)

Diabetes Mellitus

Combined surgery

Anemia ($\text{Hb} < 4.0 \text{ mmol/l}$)

Study design

Design

Study type: Observational invasive

Intervention model: Parallel

Allocation: Randomized controlled trial

Masking: Open (masking not used)

Control:	Active
Primary purpose:	Health services research

Recruitment

NL	
Recruitment status:	Recruitment stopped
Start date (anticipated):	07-04-2008
Enrollment:	279
Type:	Actual

Ethics review

Approved WMO	
Date:	08-02-2008
Application type:	First submission
Review commission:	METC Amsterdam UMC

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register	ID
CCMO	NL17356.029.07