

Effectivity of outpatient schemafocused therapy for patients with a borderline personality disorder: a pilot study

Published: 28-05-2008

Last updated: 15-05-2024

To compare the effectivity of individual SFT as described by Giesen-Bloo et al (2006) versus group SFT in patients with borderline personality disorder in an outpatient setting.

Ethical review	Approved WMO
Status	Recruiting
Health condition type	Personality disorders and disturbances in behaviour
Study type	Interventional

Summary

ID

NL-OMON31733

Source

ToetsingOnline

Brief title

Outpatient group SFT for BPD

Condition

- Personality disorders and disturbances in behaviour

Synonym

Borderline personality disorder

Research involving

Human

Sponsors and support

Primary sponsor: riagg-Maastricht

Source(s) of monetary or material Support: Ministerie van OC&W

Intervention

Keyword: BPD, group therapy, Schemafocused therapy, treatment effectivity

Outcome measures

Primary outcome

semi-structured interview: Borderline Personality Disorder Severity Index-IVth edition (BPDSI-IV). Measurements take place pre-treatment and during therapy every six months until 6 months post treatment of group therapy.

Secondary outcome

Self report questionnaires assessing quality of life (EuroQol and WhoQol), general psychopathologic dysfunction (SCL90), and specific SFT-concepts (Young Schema Questionnaire and Schema Modi Inventory-Revised).

Study description

Background summary

The present study is a continuation of a previous study on the effectivity of outpatient treatment of Borderline Personality Disorder (BPD) by means of Schema Focused Therapy (SFT) as presented by Giesen-Bloo (2006). In this study SFT proved to be effective in reducing borderline personality disorder-specific and general psychopathologic dysfunctioning and in improving quality of life. Due to long duration of the therapy, treatment is costly and waitinglists are long. Because of these reasons SFT has been implemented in group therapy in several institutions for mental health care in the Netherlands. As far as we know, the effectivity of this group-SFT has not been studied so far.

Study objective

To compare the effectivity of individual SFT as described by Giesen-Bloo et al (2006) versus group SFT in patients with borderline personality disorder in an outpatient setting.

Study design

Design: A pilotstudy about treatment effectivity: two group design in which historical data of the individual SFT are being compared with the data from the group-SFT (to be collected in the future years).

Intervention

Patients receive one group session and one individual session on a weekly basis (the individual sessions are continued as long as necessary to facilitate group sessions, these will be phased out as soon as possible) , over a period of two years.

In both sessions (group as well as individual) schema focused interventions will be applied and will be tuned by intervention of therapists on a two weekly basis.

Study burden and risks

Over a period of 30 months 6 measurements will take place, that take six hours each. Every measurement will be divided over two sessions. The total extent of the burden will be approximately 36 hours. There are no risks in attending the questionnaire assessment.

Contacts

Public

riagg-Maastricht

parallelweg 45-47
6221 BD Maastricht
NL

Scientific

riagg-Maastricht

parallelweg 45-47
6221 BD Maastricht
NL

Trial sites

Listed location countries

Netherlands

Eligibility criteria

Age

Adults (18-64 years)

Elderly (65 years and older)

Inclusion criteria

Diagnosis of Borderline Personality Disorder, by means of clinical (semi-structured) interview (Borderline personality Disorder Severity Index-IVth edition, score>20)

Speaking and writing of Dutch language

18-60 years old

Exclusion criteria

Diagnosis of mental disorders which interfere with regular treatment or group treatment: psychotic disorders, subthreshold narcissistic personality disorder or anti social personality disorder, addiction when detoxification is needed, mentally disabled persons

Study design

Design

Study phase:	2
Study type:	Interventional
Masking:	Open (masking not used)
Control:	Uncontrolled
Primary purpose:	Treatment

Recruitment

NL	
Recruitment status:	Recruiting
Start date (anticipated):	04-09-2008
Enrollment:	20
Type:	Actual

Ethics review

Approved WMO

Date: 28-05-2008

Application type: First submission

Review commission: METC academisch ziekenhuis Maastricht/Universiteit Maastricht, METC azM/UM (Maastricht)

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

ID: 26741

Source: NTR

Title:

In other registers

Register	ID
CCMO	NL20841.068.07
Other	nog niet toegekend
OMON	NL-OMON26741