

# Screening for sleep apnea using home recording of the double-loop gain as a measure of periodic breathing

Published: 05-12-2008

Last updated: 07-05-2024

Primary objectives:1. To determine the DLI threshold with optimal sensitivity and specificity. The DLI threshold is the DLI value above which the test is considered positive (and below which the test is considered negative). The optimal DLI...

<b>Ethical review</b>	Approved WMO
<b>Status</b>	Recruiting
<b>Health condition type</b>	Sleep disturbances (incl subtypes)
<b>Study type</b>	Observational non invasive

## Summary

### ID

NL-OMON31904

### Source

ToetsingOnline

### Brief title

Sleep apnea and periodic breathing

### Condition

- Sleep disturbances (incl subtypes)

### Synonym

Sleep Apnea Syndrome

### Research involving

Human

### Sponsors and support

**Primary sponsor:** Medisch Centrum Alkmaar

**Source(s) of monetary or material Support:** stichting pulmoscience

## Intervention

**Keyword:** periodic breathing, screening, sleep apnea

## Outcome measures

### Primary outcome

Area under the ROC-curve for DLI

### Secondary outcome

not applicable

## Study description

### Background summary

Sleep apnea syndrome (SAS) is characterized by repetitive events of apnea and hypopnea, usually accompanied by snoring, disturbed sleep and excessive daytime-sleepiness. These events are often part of a periodic breathing pattern, in which relative hyperventilation is followed by apnea or hypopnea. In this pattern ventilation and oxygen saturation usually oscillate with a period of 30 - 60 seconds.

Recently we described the \*double-loop gain\* of the respiratory control system as a measure of periodic breathing. This is a frequency-dependent variable which describes 1) the tendency of the respiratory system to oscillate at a given frequency and 2) the degree to which the relation between oscillations in ventilation and oscillations in arterial blood gas values is linear. The underlying hypothesis is that periodic breathing results from negative feedback regulation of arterial O<sub>2</sub> and CO<sub>2</sub> pressure through the chemoreflexes. The double-loop gain describes the gain in the negative feedback loop under the assumption that accidental changes occur in both ventilation and arterial blood gas pressures. A simple version of the double-loop gain is derived from nasal pressure changes and arterial O<sub>2</sub> saturation. From all-night recordings, the \*double-loop index\* (DLI) can be derived, which is determined by the time during which the double-loop gain exceeds a given threshold.

Currently, the presence of sleep-apnea and its clinical significance are determined by the apnea-hypopnea index (AHI), using in-hospital sleep recording. With a growing number of referrals, waiting lists for sleep registration are emerging. Screening for SAS using home-measurement of nasal pressure and SaO<sub>2</sub> seems to be a good alternative. We hypothesize that the DLI derived from these signals gives a better reflection of the pathophysiology of the disease than the AHI, which does not take into account the inherent

periodicity of the breathing pattern. As a result, we expect that the DLI improves the distinction between healthy and diseased subjects in comparison to the simple counting of apneas and hypopneas. This is reflected by a higher area under the ROC curve, which describes the sensitivity and specificity of the test.

## **Study objective**

Primary objectives:

1. To determine the DLI threshold with optimal sensitivity and specificity. The DLI threshold is the DLI value above which the test is considered positive (and below which the test is considered negative). The optimal DLI threshold will be taken as the value that gives the highest area under the ROC curve.
2. To test the hypothesis that the sensitivity and specificity of the screening are higher when the DLI is used instead of the AHI (both derived from the same home measurements of nasal pressure and SaO<sub>2</sub>).

Secondary objective:

1. To assess the repeatability of the DLI using home and in-hospital recordings of nasal pressure and SaO<sub>2</sub>.

## **Study design**

observational study without intervention

## **Study burden and risks**

In addition to standard in hospital sleep registration patients are asked to perform a one night registration of nasal pressure and O<sub>2</sub> saturation at home. They have to collect the equipment and return it the next day. No known risks are associated with this procedure. In the future this screening method may reduce the number of hospital admissions for sleep registration.

## **Contacts**

### **Public**

Medisch Centrum Alkmaar

Wilhelminalaan 12  
1815 JD Alkmaar  
Nederland

**Scientific**

Medisch Centrum Alkmaar

Wilhelminalaan 12

1815 JD Alkmaar

Nederland

## **Trial sites**

### **Listed location countries**

Netherlands

## **Eligibility criteria**

### **Age**

Adults (18-64 years)

Elderly (65 years and older)

### **Inclusion criteria**

possible sleep apnea syndrome

> 18 yr

outpatient

able and willing to use the necessary equipment to measure nasal pressure and O2 saturation at home

### **Exclusion criteria**

hospitalized patients

< 18 yr

not able to use the necessary equipment

## **Study design**

## Design

**Study type:** Observational non invasive

Masking: Open (masking not used)

Control: Uncontrolled

Primary purpose: Diagnostic

## Recruitment

NL

Recruitment status: Recruiting

Start date (anticipated): 01-08-2009

Enrollment: 50

Type: Actual

## Ethics review

Approved WMO

Date: 05-12-2008

Application type: First submission

Review commission: METC Noord-Holland (Alkmaar)

## Study registrations

### Followed up by the following (possibly more current) registration

No registrations found.

### Other (possibly less up-to-date) registrations in this register

No registrations found.

## In other registers

**Register**

CCMO

**ID**

NL21436.094.08