

Effects of 5 or 10 gram of the protein hydrolysate InsuVital* on serum insulin and glucose levels in patients with type 2 diabetes mellitus and the influence of varying carbohydrate loads

Published: 20-11-2007

Last updated: 11-05-2024

Part I: The objective of Part I of this study is the efficacy of a single 5-g and 10-g dose of InsuVital* on blood levels of insulin and glucose in patients with T2DM. Part II: The objective of Part II of this study is to assess the effect of a 10-g...

Ethical review	Approved WMO
Status	Pending
Health condition type	Glucose metabolism disorders (incl diabetes mellitus)
Study type	Interventional

Summary

ID

NL-OMON32298

Source

ToetsingOnline

Brief title

Low dose InsuVital in type 2 Diabetes Mellitus

Condition

- Glucose metabolism disorders (incl diabetes mellitus)

Synonym

diabetes mellitus type 2

Research involving

Human

Sponsors and support

Primary sponsor: DSM Food Specialties

Source(s) of monetary or material Support: DSM Food Specialties

Intervention

Keyword: insuvital, protein hydrolysate, type 2 diabetes mellitus

Outcome measures

Primary outcome

Serum concentrations and AUC of glucose and insulin.

Secondary outcome

na

Study description

Background summary

There is accumulating evidence that amino acids such as leucine play a role as insulin

secretagogues. One possible clinical application that is currently explored is a protein hydrolysate (InsuVital*). Research with this product has shown that co-ingestion of this product with carbohydrate augments the insulin response and enhances glucose disposal.

Previous experiments were carried out with a relatively high dose of protein. Hence, information on interventions with a lower protein load is necessary.

Part I of the current study will address the efficacy of 5 and 10 g of InsuVital in lowering blood levels of insulin and glucose in patients with type 2 diabetes mellitus (T2DM).

Previous experiments were also carried out with a high dose of carbohydrate. It is not known if InsuVital* is efficacious in the presence of lower carbohydrate doses.

Part II of the current study will therefore address the efficacy of a fixed dose of InsuVital, combined with either a low or a high carbohydrate load in lowering blood levels of insulin and glucose in patients with T2DM.

Study objective

Part I: The objective of Part I of this study is the efficacy of a single 5-g

and 10-g dose of InsuVital* on blood levels of insulin and glucose in patients with T2DM.

Part II: The objective of Part II of this study is to assess the effect of a 10-g dose of InsuVital* on blood levels of insulin and glucose in patients with T2DM, combined with a 25 or 50g carbohydrate load.

Study design

Randomized, placebo-controlled, double-blind, cross-over study with 3 study-days, separated by 7-day intervals.

Intervention

Part I: The treatments will consist of a drink that will be freshly prepared prior to use. The drink will be administered as a single oral bolus (300 mL) containing 50 g of carbohydrate (50% glucose and 50% maltodextrin) with 0, 5, or 10 g InsuVital*.

Part II: Patients will receive a freshly prepared drink containing 25 or 50 g of carbohydrate (50% glucose and 50% maltodextrin) with 10 g InsuVital* or 50 g without InsuVital* as a negative control.

Drinks will be flavored by adding 0.2 g sodium saccharinate, 1.8 g citric acid, and 5 g cream vanilla flavor (Quest International) per liter of beverage.

Study burden and risks

na

Contacts

Public

DSM Food Specialties

PO Box 1
2600 MA Delft
Nederland

Scientific

DSM Food Specialties

PO Box 1
2600 MA Delft
Nederland

Trial sites

Listed location countries

Netherlands

Eligibility criteria

Age

Adults (18-64 years)

Elderly (65 years and older)

Inclusion criteria

- Males or females, 18-70 years old.
- Fasting glucose level > 7 mmol/L after 2 days refraining from medication.
- Are on stable medication with biguanides for at least 3 months.
- Prepared and able to give written informed consent;

Exclusion criteria

- Use of insulin, sulfonylurea derivatives, meglitinides or other antidiabetic drugs except biguanides;
- BMI > 35 kg/m²;
- Females who are pregnant, have the intention to become pregnant within the study period, or who are lactating;
- A present and clinically significant history of ischemic heart disease (such as angina pectoris with an incidence of more than one attack/month), acute myocardial infarction within one year prior to the study or congestive heart failure (defined as NYHA class III or IV);
- Uncontrolled hypertension;
- Active, proliferative retinopathy
- Active or history of liver disease or impaired renal function (defined as a creatinin clearance calculated with the Cockcroft-Gault formula below 60 ml/min);
- Participation in a trial within 3 months prior to the start of the study or more then 4 times a year;
- Loss of 250 ml or more of blood within 3 months prior to screening;
- Any clinical condition, including use of co-medication or laboratory test results that in the opinion of the investigators may jeopardize the health status of the participants.

Study design

Design

Study type:	Interventional
Intervention model:	Crossover
Allocation:	Randomized controlled trial
Masking:	Open (masking not used)
Control:	Placebo
Primary purpose:	Treatment

Recruitment

NL	
Recruitment status:	Pending
Start date (anticipated):	01-01-2008
Enrollment:	24
Type:	Anticipated

Ethics review

Approved WMO	
Application type:	First submission
Review commission:	METC Leids Universitair Medisch Centrum (Leiden)

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register	ID
CCMO	NL19875.058.07