

# Effect of testosterone administration in Social Anxiety Disorder.

Published: 24-11-2009

Last updated: 04-05-2024

<b>Ethical review</b>	Approved WMO
<b>Status</b>	Recruitment stopped
<b>Health condition type</b>	Anxiety disorders and symptoms
<b>Study type</b>	Interventional

## Summary

### ID

NL-OMON32832

### Source

ToetsingOnline

### Brief title

Testosterone administration in SAD

### Condition

- Anxiety disorders and symptoms

### Research involving

Human

### Sponsors and support

**Primary sponsor:** Universiteit Leiden

**Source(s) of monetary or material Support:** NWO

### Intervention

**Keyword:** Social Anxiety Disorder, Social behavior, Social Fobia, Testosterone

### Outcome measures

## Study description

## Contacts

### Public

Universiteit Leiden

Wassenaarseweg 52

2300 RB Leiden

NL

### Scientific

Universiteit Leiden

Wassenaarseweg 52

2300 RB Leiden

NL

## Trial sites

### Listed location countries

Netherlands

## Eligibility criteria

### Age

Adults (18-64 years)

Elderly (65 years and older)

### Inclusion criteria

### Exclusion criteria

## Study design

## Design

Study type:	Interventional
Intervention model:	Crossover
Allocation:	Randomized controlled trial
Masking:	Double blinded (masking used)
Control:	Placebo
Primary purpose:	Other

## Recruitment

NL	
Recruitment status:	Recruitment stopped
Start date (anticipated):	21-04-2010
Enrollment:	40
Type:	Actual

## Ethics review

Approved WMO	
Date:	24-11-2009
Application type:	First submission
Review commission:	METC Leids Universitair Medisch Centrum (Leiden)

## Study registrations

### Followed up by the following (possibly more current) registration

No registrations found.

### Other (possibly less up-to-date) registrations in this register

No registrations found.

## In other registers

### Register

CCMO

### ID

NL29675.058.09