

The Santiago-study

An observational study on the vascular and metabolic effects of a pilgrimage.

Published: 21-04-2009

Last updated: 10-08-2024

In this study the effects of a two-week moderately intense-intense physical activity on vascular function, glucose-metabolism, lipid-metabolism and adipose tissue function will be analyzed.

Ethical review	Approved WMO
Status	Recruitment stopped
Health condition type	Glucose metabolism disorders (incl diabetes mellitus)
Study type	Observational invasive

Summary

ID

NL-OMON33351

Source

ToetsingOnline

Brief title

The Santiago Study

Condition

- Glucose metabolism disorders (incl diabetes mellitus)
- Vascular disorders NEC

Synonym

decreased vascular function, endothelial dysfunction

Research involving

Human

Sponsors and support

Primary sponsor: Universitair Medisch Centrum Utrecht

Source(s) of monetary or material Support: Kostenplaats vasculaire geneeskunde

Intervention

Keyword: Endothelial function, Insulin sensitivity, Physical exercise, Pilgrimage

Outcome measures

Primary outcome

Differences in FMD measurement results between the exercise group and control group directly before and after the walking-route. (ΔFMD - ΔFMD)

Secondary outcome

1. Differences in EndoPAT measurement results between the exercise group and control group directly before and after the walking-route. ($\Delta\text{EndoPAT}$ - $\Delta\text{EndoPAT}$)

2. Differences in PWV measurement results between the exercise group and control group directly before and after the walking-route. (ΔPWV - ΔPWV)

3. Differences in insulin resistance, glucose concentrations, lipid concentrations and adipokine concentrations between the exercise group and control group directly before and after the walking-route

Study description

Background summary

Rationale:

A sedentary lifestyle has been linked to increased cardiovascular morbidity and mortality. A trend towards decreased physical activity in the past decades has led to an increased prevalence of obesity and obesity-related complications such as cardiovascular disease, diabetes and cancer. It has been shown in different populations that exercise training programs lead to an improvement of the cardiovascular risk profile by enhancing insulin sensitivity, altering the lipid profile and reducing blood pressure. Endothelial dysfunction appears

early in the process of atherosclerosis and has been associated with cardiovascular disease. Endothelial function may improve after intervening in cardiovascular risk factors. Likewise, exercise has been shown to improve endothelial function in several previous studies.

In this study the effects of a two-week moderately intense-intense physical activity on vascular function, glucose-metabolism, lipid-metabolism and adipose tissue function will be analyzed in healthy participants of a Santiago pilgrimage compared to controls.

Study objective

In this study the effects of a two-week moderately intense-intense physical activity on vascular function, glucose-metabolism, lipid-metabolism and adipose tissue function will be analyzed.

Study design

Observational cohort study, with a control group.

Study burden and risks

The Santiago-study is an observational study. Non-invasive techniques will be used to assess endothelial function, vascular stiffness, central blood pressure and heart rate variability.

Invasive procedures are limited to vena punctures and finger pricks to obtain blood samples (total amount of \pm 85 ml of blood during the study per study subject).

4 Visits at the study site will be planned, which will last for approximately 1.5 hours each.

Contacts

Public

Universitair Medisch Centrum Utrecht

Heidelberglaan 100

3584 CX

Nederland

Scientific

Universitair Medisch Centrum Utrecht

Heidelberglaan 100

3584 CX

Nederland

Trial sites

Listed location countries

Netherlands

Eligibility criteria

Age

Adults (18-64 years)

Elderly (65 years and older)

Inclusion criteria

Inclusion criteria

1. Male or female subjects, 40-70 years of age
2. Voluntary participation in the Santiago de Compostela pilgrimage (only applicable for the exercise group)
3. Written and signed informed consent

Exclusion criteria

1. A history of diabetes mellitus or use of glucose-lowering medication
2. A recent history (within the last year) of alcohol or drug abuse or dependence
3. Changes in blood pressure lowering medication in the previous month and/or insufficiently controlled hypertension (systolic blood pressure >170 mmHg)
4. Usage of any lipid-lowering medication
5. A history of mental instability or major psychiatric illness not adequately controlled and stable on pharmacotherapy
6. A current (advanced) malignancy and/or current treatment for malignancy
7. A history of significant aortic coarctation
8. Significant deformity of fingers of either hand prohibiting use of the same digit on both hands for EndoPAT-measurement
9. A history of upper extremity thrombosis

Study design

Design

Study type:	Observational invasive
Intervention model:	Other
Allocation:	Non-randomized controlled trial
Masking:	Open (masking not used)

Primary purpose: Basic science

Recruitment

NL	
Recruitment status:	Recruitment stopped
Start date (anticipated):	08-05-2009
Enrollment:	60
Type:	Actual

Ethics review

Approved WMO	
Date:	21-04-2009
Application type:	First submission
Review commission:	METC Universitair Medisch Centrum Utrecht (Utrecht)

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register

CCMO

ID

NL26889.041.09