

# The effect of Neuromuscular Electrical Stimulation on post-prandial muscle protein accretion in healthy elderly men

Published: 25-07-2012

Last updated: 26-04-2024

The aim of this study is to investigate the effect of a single bout of NMES on muscle protein synthesis rates in healthy elderly males.

<b>Ethical review</b>	Approved WMO
<b>Status</b>	Recruitment stopped
<b>Health condition type</b>	Other condition
<b>Study type</b>	Interventional

## Summary

### ID

NL-OMON36866

### Source

ToetsingOnline

### Brief title

ES-PRO

### Condition

- Other condition
- Protein and amino acid metabolism disorders NEC
- Muscle disorders

### Synonym

disuse atrophy

### Health condition

muscle metabolism

### Research involving

Human

## Sponsors and support

**Primary sponsor:** Universiteit Maastricht

**Source(s) of monetary or material Support:** Ministerie van OC&W

## Intervention

**Keyword:** Elderly, Neuromuscular electrical stimulation, Protein intake, Skeletal muscle

## Outcome measures

### Primary outcome

Muscle protein synthesis rates, expressed as fractional synthetic rate (FSR).

### Secondary outcome

Muscle tracer enrichment, digestion and absorption kinetics of a casein drink influenced by NMES.

## Study description

### Background summary

With human aging there is a gradual but progressive decline in skeletal muscle mass, termed sarcopenia.. While the underlying cause of sarcopenia is likely to be multifaceted, a primary factor is that elderly individuals frequently experience short periods of muscle disuse following limb immobilization or bed-rest (due to injury or illness) causing rapid muscle loss. Elderly individuals seem to be more susceptible to muscle disuse atrophy and are less likely to fully regain their muscle tissue during subsequent rehabilitation when compared to the young. Muscle loss during a period of disuse is attributed to an impairment of muscle protein synthesis rates. Accordingly, feasible strategies for maintaining muscle protein synthesis rates in elderly individuals, and thus having the potential to attenuate the loss of muscle mass during disuse need to be pursued. Local neuromuscular electrical stimulation (NMES) offers such a potential strategy but, as yet, remains relatively unexplored.

### Study objective

The aim of this study is to investigate the effect of a single bout of NMES on

muscle protein synthesis rates in healthy elderly males.

## **Study design**

10 healthy, elderly men will consume a 20g bolus of intrinsically L[13C]phenylalanine labelled casein protein immediately following 70 min of one-legged NMES. Regular blood samples will be collected and muscle biopsies will be obtained immediately prior to protein ingestion and 240 min after ingestion from both legs to determine de novo muscle protein synthesis rates from both the stimulated (STIM) and un-stimulated control (CON) legs.

## **Intervention**

Consumption of a 20g bolus of intrinsically L[13C]phenylalanine labelled casein protein and 70 min of one-legged NMES.

## **Study burden and risks**

The risks involved in participating in this experiment are minimal. Insertion of the catheters in a vein is comparable to a normal blood draw and the only risk is a small local hematoma. This is the same for the muscle biopsies. The incision made for obtaining the muscle biopsy will be done by an experienced physician, following local anesthetics of the skin and muscle fascia, and will heal completely. The test beverages contain intrinsically labeled dietary protein which is safe for human consumption and has been used in previous studies (MEC 11-3-057 and MEC 11-3-088). NMES carries no potential risks other than slight skin irritation from the surface electrodes.

## **Contacts**

### **Public**

Universiteit Maastricht

Universiteitssingel 50 room 2.208

Maastricht 6229 ER

NL

### **Scientific**

Universiteit Maastricht

Universiteitssingel 50 room 2.208

Maastricht 6229 ER

NL

## Trial sites

### Listed location countries

Netherlands

## Eligibility criteria

### Age

Adults (18-64 years)

Elderly (65 years and older)

### Inclusion criteria

Male

Age 65 - 85 years

$18.5 < \text{BMI} < 30 \text{ kg/m}^2$

### Exclusion criteria

- Type II diabetes
- All co morbidities interacting with mobility and muscle metabolism of the lower limbs (e.g. arthrosis, arthritis, spasticity/rigidity, all neurological disorders and paralysis).
- Use of anticoagulants, blood diseases, allergy for lidocain
- Use of NSAIDs and acetylsalicylic acid
- Patients suffering from PKU (Phenylketonuria)
- Presence of implantable cardioverter defibrillator and/or pacemaker
- Performed regular resistance type exercise in the past 6 months
- Use of any tools to assist during walking (cane/ crutches/ walker)
- (Partial) foot/ leg amputation

## Study design

### Design

Study type: Interventional

Intervention model: Crossover

Allocation:	Randomized controlled trial
Masking:	Open (masking not used)
Control:	Placebo
Primary purpose:	Basic science

## Recruitment

NL	
Recruitment status:	Recruitment stopped
Start date (anticipated):	24-10-2012
Enrollment:	12
Type:	Actual

## Medical products/devices used

Generic name:	Tensmed S84 (neuromuscular electrical stimulation)
Registration:	Yes - CE intended use

## Ethics review

Approved WMO	
Date:	25-07-2012
Application type:	First submission
Review commission:	METC academisch ziekenhuis Maastricht/Universiteit Maastricht, METC azM/UM (Maastricht)
Approved WMO	
Date:	10-09-2012
Application type:	Amendment
Review commission:	METC academisch ziekenhuis Maastricht/Universiteit Maastricht, METC azM/UM (Maastricht)

## Study registrations

### Followed up by the following (possibly more current) registration

No registrations found.

**Other (possibly less up-to-date) registrations in this register**

No registrations found.

**In other registers**

Register	ID
ClinicalTrials.gov	NCT01615276
CCMO	NL40564.068.12