

Cost-effectiveness of subcutaneous immunotherapy in adults with allergic rhinitis

Published: 28-04-2009

Last updated: 19-03-2025

First, to estimate the cost-effectiveness of SCIT with tree pollen (TP), grass pollen (GP), and house dust mites (HDM) - the most prevalent allergies treated with SCIT - or combinations compared with UC. Second, to estimate the clinical efficacy of...

Ethical review	Approved WMO
Status	Recruitment stopped
Health condition type	Allergic conditions
Study type	Interventional

Summary

ID

NL-OMON37299

Source

ToetsingOnline

Brief title

AIRFORCE

Condition

- Allergic conditions

Synonym

allergic rhinitis, hay fever

Research involving

Human

Sponsors and support

Primary sponsor: Erasmus MC, Universitair Medisch Centrum Rotterdam

Source(s) of monetary or material Support: ZonMw

Intervention

Keyword: adult, allergic rhinitis, costs and cost analyses, immunotherapy

Outcome measures

Primary outcome

Cost-effectiveness: the costs per successfully treated patient, based on a global assessment of efficacy by the patient after the allergen peak exposure period in year 2 for the group that starts in 2009 and year one for the group that starts in 2010.

Clinical effectiveness: the difference in mean daily total nasal symptom scores for multi-sensitized patients in the peak exposure periods after one year.

Secondary outcome

Cost-effectiveness: the costs per symptom-free day, the costs per QALY, the costs per unit of difference between groups in the RQLQ score (disease specific quality of life).

Clinical effectiveness: the difference in mean daily total symptom scores after two years and in several subgroups; percentage of days with anti-allergic medication use, the percentage of *well days*, visual analogue scale; disease specific quality of life; global assessment; safety and adherence.

Study description

Background summary

The prevalence of allergic rhinitis is estimated at 23%. Reported annual costs of allergic rhinitis in different countries vary from \approx 1,543 to \approx 4,260 per adult. Apart from usual care (UC), which consists of symptomatic, anti-allergic medication, subcutaneous immunotherapy (SCIT) with allergens has proven

long-term effects on symptoms of both rhinitis and asthma. However SCIT is expensive and its cost-effectiveness has not been proven. Furthermore, no data are available on the efficacy of SCIT in multi-sensitized patients using more than one allergen.

Study objective

First, to estimate the cost-effectiveness of SCIT with tree pollen (TP), grass pollen (GP), and house dust mites (HDM) - the most prevalent allergies treated with SCIT - or combinations compared with UC. Second, to estimate the clinical efficacy of SCIT with a combination of two or three allergens (TP/GP/HDM) in multi-sensitized patients.

Study design

Multicenter randomized controlled open clinical trial with two parallel treatment groups

Intervention

SCIT with TP, GP, HDM or a combination plus UC or UC only, for 2 years (start autumn 2009) or 1 year (start autumn 2010).

Study burden and risks

Screening - 1 contact by phone and 1 site visit (45-60 min): questionnaire, one blood sample, physical examination (length/weight, if applicable nose inspection) and if applicable one lung function test (spirometry).

During the study - No extra site visits. Participants will be contacted by phone every 3 months (10 minutes per call). One blood sample after one year. Diary card in peak exposure period: 6-8 weeks per allergen (5-10 minutes per day). Questionnaires outside peak exposure period: approximately every 2 months (15-20 minutes per assessment).

Immunotherapy - The allergen extracts for subcutaneous administration are registered. Local side effects (itching, redness, swelling) are frequent and usually mild. Systemic allergic reactions are rare.

Contacts

Public

Erasmus MC, Universitair Medisch Centrum Rotterdam

Postbus 2040
3000 CA Rotterdam

NL

Scientific

Erasmus MC, Universitair Medisch Centrum Rotterdam

Postbus 2040

3000 CA Rotterdam

NL

Trial sites

Listed location countries

Netherlands

Eligibility criteria

Age

Adults (18-64 years)

Elderly (65 years and older)

Inclusion criteria

* 18-45 years

* Clinically relevant moderate to severe allergic rhinitis due to a sensitization for one, two or three of the following allergens: tree pollen (TP), grass pollen (GP) and/or house dust mite (HDM). For each allergen (TP, GP, HDM) the following 3 criteria are evaluated. A sensitization for an allergen is considered clinically relevant and the rhinitis moderate-severe if:

1) specific IgE ≥ 0.7 kU/l (Phadia)

2) retrospective total symptom score (RSS) ≥ 4 : participants will score 4 nose symptoms (sneezing, itching nose, watery running nose, nasal blockage) during the previous peak exposure period (TP April 1-May 15; GP May 15-June 30; HDM September 1-October 31) on a 0-3 scale (0=none, 1=mild, 2=moderate, 3=severe; maximum total score=12).

3) the presence of ≥ 1 of the following complaints due to rhinitis during the previous season: sleep disturbance; impairment of daily activities; leisure and/or sport; impairment of school or work; troublesome symptoms (QOLs).

Protocol exception: A sensitization for an allergen (specific IgE ≥ 0.7 kU/l (Phadia)) is also considered clinically relevant and the rhinitis moderate-severe if: RSS=3 and QOLs ≥ 3 or QOLs=0 and RSS ≥ 9 .

* Signed informed consent

Exclusion criteria

- * Severe/instable asthma:
 - FEV1 \leq 70% predicted and/or FEV1/FVC $<$ 70
 - Asthma exacerbation requiring prednisolon treatment, visit to a first aid station and/or hospitalisation in the preceding 12 months.
- * Specific IgE \geq 0.7 kU/l to animals the patient is in daily contact with
- * Immunotherapy in preceding 5 years
- * Anatomical disorders of the nose
- * Language barrier
- * No daily access to internet (because of web based questionnaires)
- * Contraindications to immunotherapy (according to international guidelines; i.e. history of anaphylaxis; immunosuppressive treatment etc)

Study design

Design

Study phase:	4
Study type:	Interventional
Intervention model:	Parallel
Allocation:	Randomized controlled trial
Masking:	Open (masking not used)
Control:	Active
Primary purpose:	Treatment

Recruitment

NL	
Recruitment status:	Recruitment stopped
Start date (anticipated):	05-10-2009
Enrollment:	240
Type:	Actual

Medical products/devices used

Product type:	Medicine
Registration:	Yes - NL intended use

Product type:	Medicine
Brand name:	Alutard SQ 293 Grassen-5 (RVG 16445)
Generic name:	Engels raaigras, Beemdlangbloem, Kropaar, Grote vossestaart
Registration:	Yes - NL intended use
Product type:	Medicine
Brand name:	Alutard SQ 503 (RVG 16469)
Generic name:	Huisstofmijten (Dermatophagoides pteronyssinus)
Registration:	Yes - NL intended use

Ethics review

Approved WMO	
Date:	28-04-2009
Application type:	First submission
Review commission:	METC Erasmus MC, Universitair Medisch Centrum Rotterdam (Rotterdam)
Approved WMO	
Date:	14-08-2009
Application type:	First submission
Review commission:	METC Erasmus MC, Universitair Medisch Centrum Rotterdam (Rotterdam)
Approved WMO	
Date:	14-03-2011
Application type:	Amendment
Review commission:	METC Erasmus MC, Universitair Medisch Centrum Rotterdam (Rotterdam)
Approved WMO	
Date:	17-05-2011
Application type:	Amendment
Review commission:	METC Erasmus MC, Universitair Medisch Centrum Rotterdam (Rotterdam)

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

ID: 25681

Source: Nationaal Trial Register

Title:

In other registers

Register	ID
EudraCT	EUCTR2009-011827-30-NL
CCMO	NL25370.078.09
OMON	NL-OMON25681