

Tonotopy and Modulotopy of the Auditory Cortex

Published: 27-06-2012

Last updated: 30-04-2024

The primary objective is to measure and characterize the tonotopy and modulotopy of the primary auditory cortex in normal hearing subjects. Measurements will be performed both during a normal resting state, as well as during the performance of an...

Ethical review	Approved WMO
Status	Recruitment stopped
Health condition type	Hearing disorders
Study type	Observational non invasive

Summary

ID

NL-OMON37482

Source

ToetsingOnline

Brief title

TAMAC

Condition

- Hearing disorders

Synonym

nvt

Research involving

Human

Sponsors and support

Primary sponsor: Universitair Medisch Centrum Groningen

Source(s) of monetary or material Support: Ministerie van OC&W

Intervention

Keyword: auditory, fMRI, modulotopy, tonotopy

Outcome measures

Primary outcome

Audiometric and psychometric values obtained in a diagnostic protocol, and stimulus-evoked as well as spontaneous BOLD fMRI signals in the brain.

Secondary outcome

Effect of attention (focus on various tasks) on the fMRI results of the auditory cortex.

Study description

Background summary

Sound information that is neurally encoded in the peripheral hearing organs is transmitted towards various regions in the auditory cortex. This pathway encodes acoustic attributes of sound (e.g. frequency spectrum) and its characteristics have been studied relatively well in animal models and humans (Eggermont 2001).

Previous research into the auditory cortex has shown that there are a number of cortical fields which have a tonotopical map or a modulotopical map, i.e. organization based on sound frequency or modulation frequency, respectively (Langers & van Dijk 2011, da Costa et al 2011, Talavage et al 2004, Giraud et al 2000). The aim of this study is to identify and use the various maps to distinguish between the cortical fields.

Study objective

The primary objective is to measure and characterize the tonotopy and modulotopy of the primary auditory cortex in normal hearing subjects. Measurements will be performed both during a normal resting state, as well as during the performance of an auditory and a non-auditory task, while the frequency and amplitude modulation rate of the stimuli are varied. The tasks have the aim to focus the attention of the subject either toward the auditory stimulation, for the auditory task, or to place the subjects focus on a visual

task, away from the auditory stimulation.

Study design

Exploratory study.

Study burden and risks

The clinical diagnostic tests involve several audiometric tests, and the administration of questionnaires (approx. 2 hours.). A single fMRI scanning session will take place on the same day (approx. 2 hours). None of the procedures expose the subject to known risks.

Contacts

Public

Universitair Medisch Centrum Groningen

Hanzeplein 1
Groningen 9700 RB
NL

Scientific

Universitair Medisch Centrum Groningen

Hanzeplein 1
Groningen 9700 RB
NL

Trial sites

Listed location countries

Netherlands

Eligibility criteria

Age

Adults (18-64 years)

Elderly (65 years and older)

Inclusion criteria

Normal hearing

Exclusion criteria

Hearing disorders, standard MRI-exclusion criteria (implants etc).

Study design

Design

Study type: Observational non invasive

Masking: Open (masking not used)

Control: Uncontrolled

Primary purpose: Other

Recruitment

NL

Recruitment status: Recruitment stopped

Start date (anticipated): 01-06-2012

Enrollment: 20

Type: Anticipated

Ethics review

Approved WMO

Date: 27-06-2012

Application type: First submission

Review commission: METC Universitair Medisch Centrum Groningen (Groningen)

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register

CCMO

ID

NL40057.042.12