

# Long-term effects of green tea on gut flora, fat absorption, body composition and resting energy expenditure

Published: 04-07-2012

Last updated: 01-05-2024

To investigate the long-term effects of green tea on gut flora, fat absorption, resting energy expenditure and body composition.

<b>Ethical review</b>	Approved WMO
<b>Status</b>	Recruitment stopped
<b>Health condition type</b>	Appetite and general nutritional disorders
<b>Study type</b>	Interventional

## Summary

### ID

NL-OMON37757

### Source

ToetsingOnline

### Brief title

Long-term effects of green tea

### Condition

- Appetite and general nutritional disorders

### Synonym

obesity, overweight

### Research involving

Human

### Sponsors and support

**Primary sponsor:** Universiteit Maastricht

**Source(s) of monetary or material Support:** Ministerie van OC&W

## Intervention

**Keyword:** fat absorption, green tea, gut flora, resting energy expenditure

## Outcome measures

### Primary outcome

The primary endpoint of this study is the possible change in gut flora and fat absorption.

### Secondary outcome

Secondary study parameters/endpoints are resting energy expenditure and body composition.

## Study description

### Background summary

Green tea may have positive effects for weight control and on body composition via several approaches such as a positive effect on the gut flora, a decrease in fat absorption from the intestines and an increase in resting energy expenditure.

### Study objective

To investigate the long-term effects of green tea on gut flora, fat absorption, resting energy expenditure and body composition.

### Study design

The study will be conducted in a randomized, placebo-controlled, double-blind parallel design with four groups consisting of control groups and green tea groups with normal weight subjects and obese subjects. At three time points (baseline, 6 weeks and 12 weeks) faeces are collected for analyzing the gut flora composition and fat content. Furthermore, measurements of resting energy expenditure and body composition will be conducted. Activity will be measured during three weeks (baseline, week 6 and week 12).

## Intervention

Subjects will receive either green tea or placebo in capsule form after their baseline measurement, which they have to consume three times daily for a period of twelve weeks.

### **Study burden and risks**

The study does not include any major risk for the subjects. The collection of faeces will be harmless. Anthropometric and body composition measurements will not be invasive for the subjects. Deuterium dilution has been shown to be a safe method for determining total body water and thereby body composition. Furthermore, the air in the hood is continuously regulated. Administering green tea will not form any health risk. Green tea is a natural product, which is safe in the given dose.

## **Contacts**

### **Public**

Universiteit Maastricht

Universiteitssingel 50  
Maastricht 6229 ER  
NL

### **Scientific**

Universiteit Maastricht

Universiteitssingel 50  
Maastricht 6229 ER  
NL

## **Trial sites**

### **Listed location countries**

Netherlands

## **Eligibility criteria**

### **Age**

Adults (18-64 years)

Elderly (65 years and older)

## Inclusion criteria

A total of 100 subjects (50 males and 50 females) with a BMI between 18.5-25 kg/m<sup>2</sup> and \*30 kg/m<sup>2</sup> and aged between 18-50 years will be included in the study. All subjects will be healthy, weight stable, dietary unrestrained, not using a more than moderate amount of alcohol (>10 consumptions/wk) or more than 100 mg caffeine per day, not drinking tea, not using probiotics, being weight stable (weight change < 3kg during the last 6 months), dietary unrestrained and not using antibiotics during the last 6 months. The Dutch translation of the Three Factor Eating Questionnaire (TFEQ) will be used to determine eating behaviour (10). Non-restrained eaters (<9 times factor 1), these are persons who are not consciously occupied with food and who are not caloric restricted, will be selected. Subjects will be free of medication except for oral contraceptives use in women.

## Exclusion criteria

Subjects will be excluded if they are not healthy, smoking, using a more than moderate amount of alcohol or more than 100 mg caffeine per day, drinking tea, using probiotics, not being weight stable, dietary restraint, using medication or supplements except for oral contraceptives in women, using antibiotics or if they do not meet the criteria for BMI and age. Pregnant and lactating women, and subjects with allergies for the used food items will also be excluded from participation.

## Study design

### Design

Study type:	Interventional
Intervention model:	Parallel
Allocation:	Randomized controlled trial
Masking:	Double blinded (masking used)
Control:	Placebo
Primary purpose:	Treatment

### Recruitment

NL	
Recruitment status:	Recruitment stopped
Start date (anticipated):	01-09-2012

Enrollment: 100  
Type: Actual

## Ethics review

Approved WMO  
Date: 04-07-2012  
Application type: First submission  
Review commission: METC academisch ziekenhuis Maastricht/Universiteit Maastricht, METC azM/UM (Maastricht)

## Study registrations

### Followed up by the following (possibly more current) registration

No registrations found.

### Other (possibly less up-to-date) registrations in this register

No registrations found.

### In other registers

Register	ID
ClinicalTrials.gov	NCT01556321
CCMO	NL38773.068.11