

# Investigation of the possible influence of using blood pressure lowering drugs on plasma free metanephrines concentrations.

Published: 19-03-2009

Last updated: 06-05-2024

The objective of the study is to determine whether it is necessary to take the blood pressure lowering medication (&beta;-blocker, ACE-inhibitors or diuretics) of a patient into account prior to blood sampling for pheochromocytoma diagnostics....

<b>Ethical review</b>	Approved WMO
<b>Status</b>	Recruitment stopped
<b>Health condition type</b>	Other condition
<b>Study type</b>	Observational invasive

## Summary

### ID

NL-OMON38391

### Source

ToetsingOnline

### Brief title

Influence of blood pressure lowering drugs on plasma free metanephrines.

### Condition

- Other condition
- Adrenal gland disorders

### Synonym

high blood pressure, hypertension

### Health condition

hypertensie

## Research involving

Human

## Sponsors and support

**Primary sponsor:** Universitair Medisch Centrum Groningen

**Source(s) of monetary or material Support:** Ministerie van OC&W

## Intervention

**Keyword:** antihypertensiva, hypertension, interference, plasma metanephrines

## Outcome measures

### Primary outcome

Free metanephrines (i.e. metanefrine, normetanefrine en 3-methoxytyramine) are measured in plasma.

### Secondary outcome

Nvt

## Study description

### Background summary

Since the characteristic of pheochromocytoma is the increased catecholamine (epinephrine, norepinephrine and dopamine) production in the tumor, its diagnosis is mainly based on clinical chemical results. In order to demonstrate the increased catecholamine production, free catecholamines and/or their metabolites are measured in urine and/or plasma. The analysis of the free form of 3-O-metabolized catecholamines (metanephrine, normetanephrine and 3-methoxytyramine) in plasma is the most sensitive and specific method. Therefore the measurement of plasma free metanephrines is the golden standard for the diagnosis of pheochromocytoma. This method is, as far as known, not susceptible to analytical interferences. In contrast, physiological interference could be possible for example by the use of blood pressure lowering drugs, since such medication acts on the catecholamine biochemistry. However, it is currently unknown whether the use of these drugs causes false-positive results when measuring plasma free metanephrines.

### Study objective

2 - Investigation of the possible influence of using blood pressure lowering drugs o ... 5-05-2025

The objective of the study is to determine whether it is necessary to take the blood pressure lowering medication ( $\beta$ -blocker, ACE-inhibitors or diuretics) of a patient into account prior to blood sampling for pheochromocytoma diagnostics. Therefore the influence of the use of this medication on plasma free metanephrine concentrations is established. Every patient is its own control.

Primary reserach question:

Does the use of beta-adrenoreceptor blocking drugs, angiotensine converting enzyme inhibitors or diuretics cause an increase of plasma free metanephrines which leads to false-positive pheochromocytoma diagnosis and is it therefore necessary to stop this medication prior to blood sampling?

Secondary reserach question:

Is there a difference between the use of a  $\beta$ -blocker, ACE-inhibitor or diuretics as blood pressure lowering drug?

## **Study design**

Fifty adults are asked to join the study when through screening by the general practitioner or physician untreated hypertension, peripheral oedema, palpitations or migraine was discovered. Subsequently 10 ml extra blood is drawn during routine measurements for the determination of the metanephrines basal concentrations, and the  $\beta$ -blocker, ACE-inhibitor or diuretics are prescribed. After one-three months of treatment the patient is seen for follow-up. Blood pressure is measured again and again 10 ml extra blood is drawn for the analysis of plasma free metanephrines.

Blood is sampled in sitting position by venapunction, in addition to the routine blood sampling.

Metanephrines are quantitated by isotope dilution mass spectrometry.

## **Intervention**

Blood pressure lowering drugs in the normal treatment of hypertension.

## **Study burden and risks**

During a regular visit to the general practitioner or physician, blood will be sampled twice: before treatment with a blood pressure lowering drug and after one-three months of treatment. Medical risks due to participation in the study is minimal.

## Contacts

### Public

Universitair Medisch Centrum Groningen

Hanzeplein 1 1  
Groningen 9700 RB  
NL

### Scientific

Universitair Medisch Centrum Groningen

Hanzeplein 1 1  
Groningen 9700 RB  
NL

## Trial sites

### Listed location countries

Netherlands

## Eligibility criteria

### Age

Adults (18-64 years)

Elderly (65 years and older)

### Inclusion criteria

Untreated hypertension, peripheral oedema, palpitations or migraine.

### Exclusion criteria

Treated hypertension.

Normal or low blood pressure. If there is no indication to start with  $\beta$ -blocker, ACE-inhibitors or diuretics.

## Study design

### Design

**Study type:** Observational invasive

Masking: Open (masking not used)

Control: Uncontrolled

Primary purpose: Diagnostic

### Recruitment

NL

Recruitment status: Recruitment stopped

Start date (anticipated): 06-02-2009

Enrollment: 50

Type: Actual

## Ethics review

Approved WMO

Date: 19-03-2009

Application type: First submission

Review commission: METC Universitair Medisch Centrum Groningen (Groningen)

Approved WMO

Date: 20-09-2012

Application type: Amendment

Review commission: METC Universitair Medisch Centrum Groningen (Groningen)

## Study registrations

### Followed up by the following (possibly more current) registration

No registrations found.

## Other (possibly less up-to-date) registrations in this register

No registrations found.

## In other registers

### Register

CCMO

### ID

NL24065.042.08