# The effect of fasting-induced insulin resistance on brown adipose tissue activity, non-shivering thermogenesis and skeletal muscle mitochondrial uncoupling.

Published: 07-05-2012 Last updated: 26-04-2024

The primary objectives of this study are: 1) To determine the effect of fasting-induced insulin resistance on cold-induced glucose uptake in brown adipose tissue; 2) To determine the effect of fasting on non-shivering thermogenic capacity; 3) To...

Ethical review	Approved WMO
Status	Pending
Health condition type	Other condition
Study type	Interventional

# Summary

### ID

NL-OMON39127

### Source

ToetsingOnline

#### **Brief title**

Effect of insulin resistance on brown adipose tissue activity.

# Condition

Other condition

#### Synonym

na

#### **Health condition**

Geen

1 - The effect of fasting-induced insulin resistance on brown adipose tissue activit ... 5-05-2025

#### **Research involving** Human

### Sponsors and support

**Primary sponsor:** Universiteit Maastricht Source(s) of monetary or material Support: Europese Unie

#### Intervention

**Keyword:** Brown adipose tissue, Insulin resistance, Thermogenesis

### **Outcome measures**

#### **Primary outcome**

The main study parameters are:

- Standard uptake value (SUV) of BAT: This is used as a measure of BAT glucose

uptake and BAT activity and will be measured by means of FDG-PET/CT scanning.

- Skeletal muscle mitochondrial respiration/uncoupling: This will be determined

in skeletal muscle samples using the Oroboros 2k Oxygraph instrument present in

our laboratroy.

- Energy expenditure: This will be measured by means of indirect calorimetry using a ventilated hood.

- Ex vivo skeletal muscle glucose oxidation: This will be measured to determine insulin sensitivity. A glucose oxidation assay with insulin stimulation will be performed in skeletal muscle biopsies.

#### Secondary outcome

Secondary study parameters include:

-Body temperatures: Skin temperatures will be measured by means of iButtons and core temperature will be determined by ingestion of a telemetric pill.

- Skin perfusion: This will be measured by means of LaserDoppler flowmetry.

2 - The effect of fasting-induced insulin resistance on brown adipose tissue activit ... 5-05-2025

- Blood parameters: Venous blood will be drawn by means of a catheter placed in

the antercubital vein of the underarm. By using radioimmunoassay and high

performance liquid chromatography blood parameters will be analyzed.

- UCP-1 and beta3-receptor polymorphims

# **Study description**

#### **Background summary**

Obesity and type 2 diabetes have emerged as two of the leading health problems in modern western society. So far, simple non-surgical treatment strategies for 'diabesity' are very often ineffective. A promising new target for increasing energy expenditure, and thus inducing weight loss, is facultative thermogenesis (i.e. heat production when needed, e.g. in the cold) throught the activation of brown adipose tissue (BAT). Recently, we have shown a negative correlation between BAT activity upon mild cold exposure and body mass index (BMI), suggesting that BAT might play a significant role in the development and/or sustainability of obesity. Next to obesity per se, several animal studies and retrospective human studies have suggested that diabetic status or insulin resistance, the most important physiogical hallmark of type 2 diabetes, also contribute significantly to cold-intolerance and reduced BAT activity. However, no prospective studies so far have investigated the role of insulin resistance in BAT activity upon mild cold exposure. In addition to BAT, skeletal muscle is also suggested to play a role in non-shivering thermogenesis (NST), through the process of mitochondrial uncoupling.

#### Study objective

The primary objectives of this study are: 1) To determine the effect of fasting-induced insulin resistance on cold-induced glucose uptake in brown adipose tissue; 2) To determine the effect of fasting on non-shivering thermogenic capacity; 3) To determine the effect of fasting on skeletal muscle mitochondrial uncoupling and how this is related to non-shivering thermogenesis.

#### Study design

The study will include 2 PET/CT scans for each individual, in which cold-induced BAT activity will be measured: the first scan will be performed after a normal-fed period and the second scan will be performed after a 54-h fasting period (to induce insulin resistance). Both scanning protocols consist

3 - The effect of fasting-induced insulin resistance on brown adipose tissue activit ... 5-05-2025

of a dynamic and static PET/CT scan. To investigate the role of mitochondrial uncoupling in skeletal muscle in cold-induced thermogenesis, muscle biopsies wil be taken in both conditions. Thermogenesis will be measured using indirect calorimetry and the insulative response will be determined by thermometry and skin perfusion.

#### Intervention

Fasting period of 54 hours to induce severe insulin resistance in healthy subjects. The insulin resistance is temporarily and fully reversible.

#### Study burden and risks

The risks of this experiment will be low.

The total exposure to radioactive radiation due to the administation of 18F-FDG for the PET/CT scans is 6,4 mSv for the total study. This is equal to about 3 times the total annual background radiation in the Netherlands and is considered as a low risk and fully acceptable for medical scientific research by the International Commission on Radiological Protection (<10 mSv). Thus, the risk of health damage is very limited.

# Contacts

**Public** Universiteit Maastricht

Universiteitssingel 50 Maastricht 6229 ER NL **Scientific** Universiteit Maastricht

Universiteitssingel 50 Maastricht 6229 ER NL

# **Trial sites**

### **Listed location countries**

Netherlands

# **Eligibility criteria**

#### Age

Adults (18-64 years) Elderly (65 years and older)

### **Inclusion criteria**

Healthy, lean (BMI 18-25 kg/m2) adults Age: 18-30 years Male and female For females: use of a specific oral contraceptive (Microgynon 30 or levonorgestrel/ethinylestradiol) Caucasian Sedentary (=< times/week or =< 3 hours/week sports)

# **Exclusion criteria**

Cardiovascular diseases Hypertension (systolic/diastolic blood pressure > 140/90) Hypotension (systolic/diastolic blood pressure <90/60) Use of beta-blockers Asthma or other obstructive pulmonary diseases Diabetes Mellitus type 1 and type 2 Elevated blood glucose levels (>5.6 mmol/l) Hyperthyroidism Pregnancy Participation in earlier research or medical examinations that included PET/CT scanning Radiation therapy due to medical treatment Unstable body weight (weight gain or loss >3kg in the past 2 months) Any known or suspected obstructive disease of the gastrointestinal tract, hypo motility disorders of the gastrointestinal tract or previous gastrointestinal surgery.

# Study design

# Design

**Study type:** Interventional Masking:

Open (masking not used)

Control:	Uncontrolled
Primary purpose:	Basic science

### Recruitment

NL	
Recruitment status:	Pending
Start date (anticipated):	01-05-2012
Enrollment:	18
Туре:	Anticipated

# **Ethics review**

Approved WMO	
Date:	07-05-2012
Application type:	First submission
Review commission:	METC academisch ziekenhuis Maastricht/Universiteit Maastricht, METC azM/UM (Maastricht)
Approved WMO	
Date:	11-04-2013
Application type:	Amendment
Review commission:	METC academisch ziekenhuis Maastricht/Universiteit Maastricht, METC azM/UM (Maastricht)

# **Study registrations**

# Followed up by the following (possibly more current) registration

No registrations found.

# Other (possibly less up-to-date) registrations in this register

No registrations found.

# In other registers

Register	ID
Other	Nederlands Trialregister: TC = 3523
ССМО	NL39816.068.12