

# Posttraumatic osteoarthritis following distal radius fractures in young patients and the correlation with subjective and objective outcome measures

Published: 23-07-2013

Last updated: 15-05-2024

To determine the incidence of posttraumatic osteoarthritis following distal radius fractures in a cohort of young non-osteoporotic patients and correlation with objective and subjective outcome measures.

<b>Ethical review</b>	Approved WMO
<b>Status</b>	Recruitment stopped
<b>Health condition type</b>	Fractures
<b>Study type</b>	Observational invasive

## Summary

### ID

NL-OMON39658

### Source

ToetsingOnline

### Brief title

Osteoarthritis following distal radius fractures in young patients

### Condition

- Fractures
- Bone and joint therapeutic procedures

### Synonym

distal radius fracture, wrist fracture

### Research involving

Human

## Sponsors and support

**Primary sponsor:** Universitair Medisch Centrum Groningen

**Source(s) of monetary or material Support:** Ministerie van OC&W, Fonds de Gavere

## Intervention

**Keyword:** Fracture, Osteoarthritis, Radius, Subjective

## Outcome measures

### Primary outcome

The incidence of posttraumatic osteoarthritis.

### Secondary outcome

Subjective outcome measures objectivied with the use of validated questionnaires (PRWE, DASH, SF-36 and MHQ). Objective outcomes measured with functional tests such as range of motion and gripstrength. Radiological findings measured with X-rays.

## Study description

### Background summary

The development of posttraumatic osteoarthritis (PA) following distal radius fractures (DRFs) has been commonly described. Direct and indirect joint impact loading, soft tissue injuries, joint dislocation and intra-articular fractures, increase the risk of progressive joint degeneration that causes PA. It is thought posttraumatic osteoarthritis develops less in younger patients. However, it might be more invalidating for a young non-osteoporotic patient to develop posttraumatic osteoarthritis and loss of function following a distal radius fracture than for an older patient. The extend of the loss of function can be objectivied using functional measures, such as range of motion and grip strength. Subjective measures to objectivy loss of function as experienced by the patient can be performed using validated questionnaires. In this study, the incidence of posttraumatic osteoarthritis following a distal radius fracture is in young patients is determined. Also, the question arises what the correlation between the existence of post traumatic osteoarthritis and the objective and subjective outcome measures is following a distal radius fracture in young

patients.

## **Study objective**

To determine the incidence of posttraumatic osteoarthritis following distal radius fractures in a cohort of young non-osteoporotic patients and correlation with objective and subjective outcome measures.

## **Study design**

Descriptive cohort study

## **Study burden and risks**

According to the international commission on radiological protection (ICRP) the estimated risk for the radiological procedure (4 x-rays per subject) is categorized as IIa, very low level of risk of radiogenic cancer,  $>0.2$  and  $<2$  mSv. In comparison to the \* Rijks Instituut Voor Milieuhygiëne (RIVM)\*, the annual radiation dose in the Netherlands is 1.7 mSv. The number of visits is once for the functional physical tests combined with the radiological procedure. The physical discomfort during the functional tests and the radiological investigation is minimal. The subject will be asked to sit still for the radiological investigations for 4 times less than 1 minute.

## **Contacts**

### **Public**

Universitair Medisch Centrum Groningen

Hanzeplein 1  
Groningen 9713GZ  
NL

### **Scientific**

Universitair Medisch Centrum Groningen

Hanzeplein 1  
Groningen 9713GZ  
NL

## **Trial sites**

## Listed location countries

Netherlands

## Eligibility criteria

### Age

Adults (18-64 years)

Elderly (65 years and older)

### Inclusion criteria

All patients treated in the period 2005 until 2011 in the Medical Center Leeuwarden with a distal radius fracture of which the x-ray after the trauma and after 6 weeks is present in the archives of the Radiology Department. Men aged between the age 18-50 years and women between the age of 18-40 years at the time of presentation.

### Exclusion criteria

Fractures treated more than 7 days following initial trauma, open fractures and preexistent osteoarthritis.

## Study design

### Design

**Study type:** Observational invasive

Masking: Open (masking not used)

Control: Uncontrolled

Primary purpose: Diagnostic

### Recruitment

NL

Recruitment status: Recruitment stopped

Start date (anticipated): 01-09-2013

Enrollment: 80

Type:

Actual

## Ethics review

Approved WMO

Date:

23-07-2013

Application type:

First submission

Review commission:

RTPO, Regionale Toetsingscie Patientgebonden Onderzoek  
(Leeuwarden)

## Study registrations

### Followed up by the following (possibly more current) registration

No registrations found.

### Other (possibly less up-to-date) registrations in this register

ID: 22608

Source: NTR

Title:

### In other registers

Register	ID
Other	Aangemeld bij NTR
CCMO	NL41587.099.13
OMON	NL-OMON22608