# Longterm Skeletal Stability of Mandibular Distraction Osteogenesis

Published: 21-01-2014 Last updated: 16-11-2024

To investigate the long term stability (> 5 yrs) in patients treated with mandibular distraction osteogenesis (MDO) for correction of a Class II-occlusion. Hypotheses:1) MDO is a technique for mandibular lengthening with comparable stability to...

Ethical review	Approved WMO
Status	Will not start
Health condition type	Other condition
Study type	Observational invasive

# Summary

#### ID

NL-OMON39925

**Source** ToetsingOnline

#### **Brief title**

Longterm Skeletal Stability of Mandibular Distraction Osteogenesis

### Condition

- Other condition
- Bone disorders (excl congenital and fractures)
- · Bone and joint therapeutic procedures

#### Synonym

mandibular hypoplasia, overbite/overjet

#### **Health condition**

evaluatie stabiliteit van chirurgische interventie

#### **Research involving**

Human

### **Sponsors and support**

**Primary sponsor:** Vrije Universiteit Medisch Centrum **Source(s) of monetary or material Support:** Ministerie van OC&W

#### Intervention

Keyword: Distraction osteogenesis, Mandible, Relapse, Stability

#### **Outcome measures**

#### **Primary outcome**

Main study parameters

Cephalometric analysis will be carried out on the pre-operative,

post-distraction and these radiographs. The following measurements will be

made:

- \* SNB-angle: sella/nasion-mandibular point B
- \* SN-MP: sella/nasion- mandibular plane
- \* Point B in respect to x and y-axis
- \* x-axis= SN, y-axis: perpendicular line to x-axis

Primary parameter is to establish horizontal mandibular relapse. This is

defined as change of position of point B in a horizontal (mandibular) plane.

#### Secondary outcome

N/A

# **Study description**

#### **Background summary**

Mandibular distraction osteogenesis (MDO) is an alternative technique for mandibular lengthening to the conventional bilateral sagittal split osteotomy (BSSO) in patients with a mandibular hypoplasia. Both methods are surgical

procedures that are performed in a combined surgical-orthodontic treatment plan. Longterm skeletal stability after BSSO has been studied en reported in the literature for the conventional method. Influencing factors as the amount of advancement, type of fixation, mandibular plane (high- and low angle patients), surgical skills and bone, muscular and soft tissue have been reported. An advantage of MDO is the possibility to achieve a larger amount of lengthening (>10mm), which is required in some patients with an severe mandibular hypoplasia, e.g. syndromic patients or children. Therefore, the chances of skeletal relapse might be increased in these patients. Short- and middleterm stability of MDO in comparison to the BSSO-technique shows similar skeletal stability. However, to our knowledge the longterm stability (at least more than 5 years) of MDO has not been studied or reported in the literature. Since this technique has become a more routine procedure in orthognatic surgery, it is of importance to establish the longterm stability and amounts of possible skeletal relapse. These results could lead to recommendations to the surgical protocol and could be beneficial to future patients.

#### Study objective

To investigate the long term stability (> 5 yrs) in patients treated with mandibular distraction osteogenesis (MDO) for correction of a Class II-occlusion.

Hypotheses:

1) MDO is a technique for mandibular lengthening with comparable stability to other commonly used techniques (BSSO)

2) High-angle patients, lengthening > 10 mm and female patients (>38°) are more at risk for relapse than low-angle patients, lengthening < 10mm and male patients

#### Study design

This is a retrospective, cephalometric study to determine skeletal stability of mandibular distraction performed between 1999-2006. Patients will be recalled for a cephalometric radiograph. Cephalometric tracing will be carried out using advanced software and manual tracing by two investigators.

#### Study burden and risks

In our opinion the nature and extrent of burden and associated risk is minimal. The longterm stability (at least more than 5 years) of mandibular distraction osteogenesis has not been studied or reported in the literature. Since this technique has become a more routine procedure in orthognatic surgery, it is of importance to establish the longterm stability and amounts of possible skeletal relapse. These results could lead to recommendations to the surgical protocol and could be beneficial to future patients.

# Contacts

#### Public

Vrije Universiteit Medisch Centrum

De Boelelan 1118 Amsterdam 1087 HV NL **Scientific** Vrije Universiteit Medisch Centrum

De Boelelan 1118 Amsterdam 1087 HV NL

# **Trial sites**

### **Listed location countries**

Netherlands

# **Eligibility criteria**

#### Age

Adults (18-64 years) Elderly (65 years and older)

### **Inclusion criteria**

Mandibular hypoplasia treated with mandibular distraction osteogenesis

### **Exclusion criteria**

Follow-uptime shorter <5 years Other orthognatic surgery (genioplasty, sagittal split osteotomy)

# Study design

### Design

Study type: Observational invasive		
Masking:	Open (masking not used)	
Control:	Uncontrolled	
Primary purpose:	Treatment	

### Recruitment

NL	
Recruitment status:	Will not start
Start date (anticipated):	01-03-2014
Enrollment:	55
Туре:	Anticipated

# **Ethics review**

Approved WMO	
Date:	21-01-2014
Application type:	First submission
Review commission:	METC Amsterdam UMC

# **Study registrations**

### Followed up by the following (possibly more current) registration

No registrations found.

### Other (possibly less up-to-date) registrations in this register

No registrations found.

# In other registers

### Register

ССМО

**ID** NL40744.029.14