

Upper eyelid blepharoplasty investigating 3D and functional outcomes

Published: 17-11-2017

Last updated: 12-04-2024

The aim of this study is to objectively assess 3D and functional outcomes of upper blepharoplasty on periorbital volume, dry eyes, peripheral vision, electrical activity of the frontal muscles and wellbeing in two different surgical techniques.

Ethical review	Approved WMO
Status	Recruiting
Health condition type	Other condition
Study type	Interventional

Summary

ID

NL-OMON44420

Source

ToetsingOnline

Brief title

Upper eyelid blepharoplasty investigating 3D and functional outcomes

Condition

- Other condition

Synonym

blepharochalasis, excess skin upper eyelid

Health condition

blefarochalasis, huidoverschot van de bovenoogleden

Research involving

Human

Sponsors and support

Primary sponsor: Universitair Medisch Centrum Groningen

Source(s) of monetary or material Support: Ministerie van OC&W

Intervention

Keyword: Blepharoplasty, Three-dimensional imaging

Outcome measures

Primary outcome

The primary endpoint is orbital change in volume.

Secondary outcome

Secondary outcomes are electrical activity of the frontal and orbicularis oculi muscles, peripheral vision, presence and severity of dry eyes, quality of life and wellbeing.

Study description

Background summary

Blepharoplasty of the upper eyelids is one of the most commonly performed surgical procedures in aesthetic surgery. This surgery may be performed for aesthetic reasons, but also for functional indications. Based on the existing literature, the outcomes of an upper blepharoplasty might depend on the peri-orbital volume and the surgical technique applied. The traditional technique of upper blepharoplasty consists of the resection of skin and a strip of the orbicularis oculi muscle, and coagulation of fat. A common alternative technique involves the removal of skin only (alternative). The outcomes (esthetic and functional) between these techniques may vary. Both techniques are widely used in daily practice. However, there is yet no consensus about which technique is preferable.

Study objective

The aim of this study is to objectively assess 3D and functional outcomes of upper blepharoplasty on periorbital volume, dry eyes, peripheral vision, electrical activity of the frontal muscles and wellbeing in two different

surgical techniques.

Study design

The study design is a clinical trial. Patients are randomized in the *traditional* (resection of skin and strip of the orbicularis oculi muscle) or *alternative* (skin only) blepharoplasty treatment group. The investigators and patients are blinded during the study. Patients undergo a series of functional tests and 3D photography pre- and postoperatively. The follow up is 12 months.

Intervention

Patients undergo a blepharoplasty.

Study burden and risks

Both surgical techniques are common techniques for a upper blepharoplasty, widely used and considered not to be harmful. Specific for this study, the patient have to undergo a series of non-invasive tests. Completing these tests cost the patients 45 minutes extra per visit. During the one year follow-up, the participants have to visit the clinic 4 times of which 2 visits are combined their regular control appointments (standard care).

Contacts

Public

Universitair Medisch Centrum Groningen

Hanzeplein 1
Groningen 9700 RB
NL

Scientific

Universitair Medisch Centrum Groningen

Hanzeplein 1
Groningen 9700 RB
NL

Trial sites

Listed location countries

Netherlands

Eligibility criteria

Age

Adults (18-64 years)

Elderly (65 years and older)

Inclusion criteria

- Blepharochalasis of both eyes
- Age between 30 and 60
- Caucasian
- Both male and female patients are included
- Correction of vision between -6 and +6
- Vision of 0.5 or better (with correction)
- Fluent in Dutch
- Legally capable

Exclusion criteria

- Protruding medial fat compartment
- Ocular trauma in past
- Trauma of the orbital region in the past
- A medical history of eyelid surgery or surgery in the region of the eyebrows.
- Any current ophthalmic disease that could interfere with the ophthalmic tests
- Horner syndrome
- Blepharoptosis
- Graves* disease
- Myasthenia Gravis
- Recent (in the past 6 months) botulin toxin treatment in the upper face
- Nasolacrimal duct obstruction
- Epilepsy
- Retinal defects/diseases that cause visual field defects

Study design

Design

Study type: Interventional

Masking: Double blinded (masking used)

Control: Uncontrolled

Primary purpose: Other

Recruitment

NL

Recruitment status: Recruiting

Start date (anticipated): 20-02-2018

Enrollment: 54

Type: Actual

Ethics review

Approved WMO

Date: 17-11-2017

Application type: First submission

Review commission: METC Universitair Medisch Centrum Groningen (Groningen)

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register

CCMO

ID

NL62635.042.17