

# A randomised comparison of Eye Movement Desensitization and Reprocessing (EMDR), Narrative Exposure Therapy (KIDNET) and waiting list control in traumatized refugee children.

Published: 17-02-2014

Last updated: 19-03-2025

The study aims to answer the following research questions:1. How high/low are levels of symptomatology of depression and/or PTSD in traumatised refugee children in ASC?2. Do symptom levels decrease after a short-term trauma-focused intervention?3....

|                              |                                |
|------------------------------|--------------------------------|
| <b>Ethical review</b>        | Approved WMO                   |
| <b>Status</b>                | Recruiting                     |
| <b>Health condition type</b> | Anxiety disorders and symptoms |
| <b>Study type</b>            | Interventional                 |

## Summary

### ID

NL-OMON44793

### Source

ToetsingOnline

### Brief title

Effects of EMDR and KIDNET in refugeechildren

### Condition

- Anxiety disorders and symptoms
- Lifestyle issues

### Synonym

angst, Posttraumatic Stress Disorder (PTSD)

## Research involving

Human

## Sponsors and support

**Primary sponsor:** Stichting Centrum 45 (Oegstgeest)

**Source(s) of monetary or material Support:** gedeeltelijke door subsidie van EMDR Europe en Vereniging EMDR Nederland (VEN)

## Intervention

**Keyword:** psychotrauma, RCT, refugeechildren, Traumafocused intervention

## Outcome measures

### Primary outcome

Key variables of this study are:

1. Posttraumatic stress symptoms (disorder)
2. Quality of life
3. General non-trauma specific stress reactions (such as bedwetting, eating problems)
4. Behavioral and emotional symptoms

### Secondary outcome

Drop-out, number of sessions, number of weeks, duration of intervention, anxiety in parent(s).

## Study description

### Background summary

Traumatized refugee children present a complex constellation of symptoms. Many of them have been exposed to multiple stressful experiences, such as violence, separation and migration, and are facing difficult living circumstances. Few efficacy studies of treatments to this vulnerable group of children are available (Henley & Robinson, 2011). The evidence for the use of the

psychotherapeutic intervention method EMDR in refugee children is still in its infancy (Oras, De Ezpeleta & Ahmad, 2004) and limited when it comes to KIDNET (Onyut, et al., 2005; Catani et al., 2009; Ruf et al., 2010). On the basis of meta-analytic results, it is recommended to further explore trauma focused interventions\* incremental efficacy for refugee children in controlled studies (Rodenburg et al., 2009). This project is a collaborative effort to study the effects of EMDR and KIDNET as trauma-focused interventions for refugee children in a RCT. Its significance is related to the limited knowledge that exists in this field, with the theoretical gain that is expected, and with the relevance because of a large and growing number of children in comparable circumstances worldwide.

This study combines expertise in applying EMDR and KIDNET, in children (Rodenburgh et al., 2009; De Roos et al., 2011; Neuner et al., 2008; Onyut et al., 2005; Wanders et al., 2008) with expertise in working with this vulnerable group. Effective interventions for mental health problems related to exposure to traumatic events in children is lagging behind in comparison to adults. Additionally, it is known that children may be burdened by psycho-traumatic symptoms and hampered in their development (Van Ee, Kleber & Mooren, 2012). Children of refugees and asylum-seekers offer an extra challenge for mental health services. While violence and loss have frequently disrupted their lives, they are also subjected to changing social surroundings due to flight, frequent allocations and an uncertain future. At the same time, they have caregivers undergoing similar levels or higher levels of stress causing various mental health problems, which are thought to intertwine with mental health problems of the child through limited parental availability and their rearing practices. Complex posttraumatic disturbances may follow cumulative and chronic disrupting life-events, such as war and subsequent migration and consist of severe chronic symptoms as well as changes within the personality (Herman, 1993). The impact on further development, including their ability to integrate, can be dramatic. Therefore, it is important to offer interventions timely, efficiently and effectively. Long-term consequences need to be prevented. This study aims at optimizing their chances for recovery.

The theoretical questions that will be answered in this project are related to the importance of traumatic memories and intrusions in children with more complex symptoms of PTSD following traumatic experiences. Is trauma-focused therapy (EMDR versus KIDNET) effective in alleviating and reducing PTSD? Is this true for different age-groups: 6-8 and 12-18 years? Which factors are predictive for treatment outcomes? Outcomes of this project will help us understand how traumatic stress symptoms can be reduced in a high-risk group of children. This project is a step towards disentangling these issues. Lastly, the children included in this study are currently living in asylum centers and awaiting the decision on their asylum application. A growing number of refugees live in Western Countries, often in uncertainty of their future for longer periods of time. This study can contribute to the discussion in the mental health field if trauma focused therapy can be offered to this group as

they experience this uncertainty and live in non-optimal circumstances. Ultimately, this study can open doors for a large group of children in need, too often \*unseen\* in mental health institutions because of the lack of understanding what trauma therapy can safely offer them in this period of their lives.

## **Study objective**

The study aims to answer the following research questions:

1. How high/low are levels of symptomatology of depression and/or PTSD in traumatised refugee children in ASC?
2. Do symptom levels decrease after a short-term trauma-focused intervention?
3. Are EMDR and/or KIDNET effective in decreasing symptoms at the end of treatment and/or follow-up?
4. Are EMDR and/or KIDNET effective in improving quality of life at the end of treatment and/or follow-up?

The benefits of this project are several and related to implementation of trauma-focused methods, examining systematically effects of these types of interventions in high-risk groups and lastly dissemination of knowledge and expertise (by publication and presentation). This project will result in:

- \* starting clinical research expertise in the field of treatment of traumatized refugee children.
- \* disseminating of findings to scientific and clinical professionals (through publications, presentations).
- \* examining the merits of care as usual for this population and registration of specific type of interventions.

## **Study design**

The design of the study is a randomized controlled trial using standardized questionnaires as well as structured interviews and performance data (quality of life); children between 8 and 18 years of age will be randomly assigned to either EMDR, KIDNET or waitlist control (N=30-30-30). They will be stratified in order to obtain two equal sub-age-groups (8-12; 13-18 years). Sample size has been based on power calculations incorporating small to moderate effect sizes and risk of drop-out (software G\*power (Faul, Erdfelder, Lang, Buchner, 2007)).

Children will be assessed at intake (before first treatment session), after the last session (maximum of 8 sessions of 1 hour) and at follow-up (3 months after the last session). Moreover, participants will fill out the CRIES (Children's Revised Impact of Event Scale) each session to monitor changes closely. In the period between the last treatment session and follow-up no prolonged treatment

will be offered, unless really needed. The EMDR therapy will be performed by therapists who are trained at EMDR level-II and in providing KIDNET. They will receive monthly supervision by a registered supervisor. Treatment fidelity will be checked to assess treatment adherence in both conditions.

## **Intervention**

Participants of the study will be informed about the study and then randomly assigned to one of two interventions versus a (temporarily) wait-list condition: 1. Eye Movement Desensitization Reprocessing (EMDR), 2. Narrative Exposure Therapy (KIDNET), versus 3. Waiting list.

### **Interventions**

Both research conditions will consist of 8 weekly sessions. The EMDR treatment condition starts with one preparatory session to establish a working alliance, conduct a case conceptualisation and to obtain agreement on treatment targets.

#### **(1) EMDR**

The trauma-focused EMDR condition continues with 7 sessions aimed at reducing disturbance associated with the most distressing traumatic memories, following the Dutch version of the EMDR protocol for children and adolescents (de Roos, Beer, De Jongh & ten Broeke; based on De Jongh & Ten Broeke, 2003; Shapiro, 2001). EMDR sessions last 75 minutes.

#### **(3) KIDNET**

The trauma-focused KIDNET condition continues with 7 sessions aimed at reducing disturbance associated with the most distressing traumatic memories, in chronological order (Neuner et al., 2008; Schauer et al., 2017). Kidney sessions last 75 minutes (same amount as EMDR).

#### **(3) Waiting list control condition (WL)**

Once trauma-focused treatment has been indicated, a child may be assigned to the waiting list. This implies that a child will have to wait 8 weeks before the start of EMDR or KIDNET (second randomization).

### **Parental guidance**

In all conditions parent guidance will be offered \* with a maximum of 4 sessions 1 hour, commonly with interpreter). There is consensus in the clinical field that parent guidance in the treatment of refugee children is needed. Psychoeducation is the dominant subject of the encounters with parents (e.g., information about the interventions; what kind of reactions can be expected in their children). The interventions (in terms of duration, kind and frequency) that parents receive will be registered.

In the treatment of children there is great reluctance to prescribe medication; during this study it is therefore not to be expected that any medication will be needed as extra intervention. In the case of a

crisis, a child-psychiatrist is available to be consulted \* as is a normal procedure during treatment. Extra interventions will be noted (in the rare occasion medication will be prescribed, the child will be excluded from the study).

### **Study burden and risks**

No particular risks are being foreseen for participants. Data will be handled confidentially and coded. A subject identification code will be used to link the research data to the subject. The key to the code will be kept separate by the investigator. The handling of personal data will be according to the Dutch Personal Data Protection Act (in Dutch: De Wet Bescherming Persoonsgegevens, Wbp).

Children will be compensated for their participation with a coupon (euro 15) per measurement after treatment (M2 and M3).

## **Contacts**

### **Public**

Stichting Centrum 45 (Oegstgeest)

Rijnzichtweg 35  
Oegstgeest 2342 AX  
NL

### **Scientific**

Stichting Centrum 45 (Oegstgeest)

Rijnzichtweg 35  
Oegstgeest 2342 AX  
NL

## **Trial sites**

### **Listed location countries**

Netherlands

## **Eligibility criteria**

### **Age**

Adolescents (12-15 years)  
Adolescents (16-17 years)  
Children (2-11 years)

## Inclusion criteria

- aged 8 - 18 years; 8-12 ( $n \leq 15$ ) and 13-18 years ( $n \leq 15$ ) per group, per condition (EMDR, KIDNET, WL)
- PTSD
- speaking Dutch

## Exclusion criteria

- estimated low intelligence-level;
- severe and acute psychiatric conditions, such as high-risk suicidality, psychosis, aggressive behaviour;
- brain damage.

# Study design

## Design

|                     |                               |
|---------------------|-------------------------------|
| Study type:         | Interventional                |
| Intervention model: | Parallel                      |
| Allocation:         | Randomized controlled trial   |
| Masking:            | Single blinded (masking used) |
| Control:            | Active                        |
| Primary purpose:    | Treatment                     |

## Recruitment

|                           |            |
|---------------------------|------------|
| NL                        |            |
| Recruitment status:       | Recruiting |
| Start date (anticipated): | 05-01-2015 |
| Enrollment:               | 30         |
| Type:                     | Actual     |

## Ethics review

Approved WMO

Date: 17-02-2014

Application type: First submission

Review commission: METC Leiden-Den Haag-Delft (Leiden)

metc-ldd@lumc.nl

Approved WMO

Date: 16-06-2017

Application type: Amendment

Review commission: METC Leiden-Den Haag-Delft (Leiden)

metc-ldd@lumc.nl

## Study registrations

### Followed up by the following (possibly more current) registration

No registrations found.

### Other (possibly less up-to-date) registrations in this register

ID: 22679

Source: NTR

Title:

### In other registers

| Register | ID             |
|----------|----------------|
| CCMO     | NL40769.058.13 |
| OMON     | NL-OMON22679   |