

Experience sampling of depression, religion/spirituality and suicidality: a new stage.

Published: 02-11-2018

Last updated: 11-04-2024

To (1) evaluate the adequacy of ESM-variables and -method in measuring depression, R/S and suicidality, (2) gain insight into the variability and interrelations of depression, R/S and suicidality, and validate ESM-parameters on R/S, (3) raise...

Ethical review	Approved WMO
Status	Recruitment stopped
Health condition type	Mood disorders and disturbances NEC
Study type	Observational non invasive

Summary

ID

NL-OMON46063

Source

ToetsingOnline

Brief title

Depression and meaning-making: a new phase.

Condition

- Mood disorders and disturbances NEC

Synonym

depression

Research involving

Human

Sponsors and support

Primary sponsor: Kennisinstituut christelijke ggz

Source(s) of monetary or material Support: Stichting tot Steun van de VCVGZ.

Intervention

Keyword: depression, experience sampling, religion, suicidality

Outcome measures

Primary outcome

6 days of smartphone-based assessment, randomly distributed over the day in 10 time-slots (8:00 till 20:00), preceded and followed by several regular questionnaires.

Secondary outcome

Not applicable.

Study description

Background summary

Evidence is growing that religion and spirituality (R/S) have a protective effect on suicide and suicidality. Nevertheless, it is not clear what accounts for this effect. The longitudinal research needed to clarify this effect is scarce. The Experience Sampling Method (ESM) would be effective for studying depression, R/S and suicidality at both a within-subject and between-subject level, but such research has never been performed before.

Study objective

To (1) evaluate the adequacy of ESM-variables and -method in measuring depression, R/S and suicidality, (2) gain insight into the variability and interrelations of depression, R/S and suicidality, and validate ESM-parameters on R/S, (3) raise hypotheses and a model for further more detailed research and (4) ultimately inspire new opportunities for interventions and treatment.

Study design

An observational longitudinal study using ESM consisting of a user co-creation phase and an exploratory study. This study will be followed by a main study on the same variables.

Study burden and risks

This study is only an observational study. The risk of a raise in suicidality by this assessment is very low, a decrease of suicidality is even more probable. Suicidality is monitored both by the health care provider(s) and (assistant-)researcher. A crisis-plan is available. A subject can withdraw from the study at any moment.

Contacts

Public

Kennisinstituut christelijke ggz

Printerweg 21
AMERSFOORT 3821 AP
NL

Scientific

Kennisinstituut christelijke ggz

Printerweg 21
AMERSFOORT 3821 AP
NL

Trial sites

Listed location countries

Netherlands

Eligibility criteria

Age

Adults (18-64 years)
Elderly (65 years and older)

Inclusion criteria

Adult persons with a depressive disorder, actually treated in a specialized mental health care institution (sggz) institution.

Exclusion criteria

Persons not being able to understand or speak Dutch, with general learning disability, or not able to handle a smartphone or a tablet.

Study design

Design

Study type: Observational non invasive

Masking: Open (masking not used)

Control: Uncontrolled

Primary purpose: Basic science

Recruitment

NL

Recruitment status: Recruitment stopped

Start date (anticipated): 01-03-2019

Enrollment: 30

Type: Actual

Ethics review

Approved WMO

Date: 02-11-2018

Application type: First submission

Review commission: METC Universitair Medisch Centrum Groningen (Groningen)

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register	ID
CCMO	NL65565.042.18