

Diagnostic imaging of synovitis in knee osteoarthritis (DISKO)

Published: 30-06-2016

Last updated: 16-04-2024

Our objectives are to assess correlations and relationships between synovitis visualized on non-contrast DESS MRI and: 1) synovitis imaged on CE-MRI; 2) different inflammatory subtypes in OA; 3) other MRI features of OA; 4) knee OA symptoms and 5)...

Ethical review	Approved WMO
Status	Recruiting
Health condition type	Synovial and bursal disorders
Study type	Observational invasive

Summary

ID

NL-OMON46234

Source

ToetsingOnline

Brief title

Diagnostic imaging of synovitis in knee osteoarthritis (DISKO)

Condition

- Synovial and bursal disorders

Synonym

Synovitis; mucosal inflammation

Research involving

Human

Sponsors and support

Primary sponsor: Erasmus MC, Universitair Medisch Centrum Rotterdam

Source(s) of monetary or material Support: Ministerie van OC&W, Reumafonds

Intervention

Keyword: Knee, MRI, osteoarthritis, synovitis

Outcome measures

Primary outcome

Correlations between synovitis visualized on non-contrast DESS and: 1) CE-MRI; 2) Tissue markers of synovial inflammation and specific inflammatory subtypes; 3) Other OA features on MRI; 4) knee OA symptoms and 5) (CE)US findings related to synovitis.

Secondary outcome

1. Synovial Tissue and synovial fluid markers of synovial inflammation and specific inflammatory subtypes;
2. Other OA features on MRI, particularly compositional and morphological change in cartilage and meniscus, altered blood perfusion in synovium and subchondral bone, and osteophyte formation;
3. Knee OA symptoms, especially those related to synovitis (i.e. joint stiffness, self-reported feeling of swelling, pain patterns (constant versus intermittent pain)), recorded with validated standardized questionnaires (like ICOAP, KOOS, VAS and WOMAC).
4. US findings related to synovitis

Study description

Background summary

As the prominent role of synovitis in osteoarthritis (OA) and importance of identifying patients with synovitis for targeted treatment are increasingly

recognized, there is growing interest in imaging of synovitis in OA. Contrast-enhanced magnetic resonance imaging (CE-MRI) is the reference standard for visualizing synovitis, but incurs high costs, long scan time and potential health issues in high-risk patients, associated with the contrast agent. A promising recent innovation in MRI of synovitis is rapid diffusion weighted imaging with dual echo steady state (DESS) without the need for a contrast agent. We hypothesize that synovitis imaged with non-contrast DESS MRI shows high correlation with 1) synovitis imaged on CE-MRI; 2) different inflammatory subtypes in OA; 3) other MRI features of OA, 4) knee OA symptoms and 5) synovitis imaged with contrast enhanced ultrasound (CEUS), which is another option to visualize synovitis.

Study objective

Our objectives are to assess correlations and relationships between synovitis visualized on non-contrast DESS MRI and: 1) synovitis imaged on CE-MRI; 2) different inflammatory subtypes in OA; 3) other MRI features of OA; 4) knee OA symptoms and 5) synovitis imaged with (CE)US.

Study design

Cross-sectional prospective observational study.

Study burden and risks

Two MRI sessions, each maximal 45 minutes, and exposure to acoustic noise. No additional risk. No benefit for the patient. There is a very small risk for an allergic reaction when MRI contrast agent is administered. In patients with chronic severe kidney insufficiency (glomerular filtration rate (GFR) < 60 mL/min/1.73m²) the use of this contrast is contraindicated. In order to diminish this risk, patients will be screened before inclusion. Thereby, medication to counteract the reaction is present. The burden for the subject will be minimal. Except draining a vial of blood, administration of contrast fluid, the burden will mainly exist of time for filling out a questionnaire, undergoing a physical examination, cycling for 10 minutes on a home trainer and MRI acquisition. There will be no direct benefits for the participants in this study. All patients will receive the usual treatment from their treating physician. The patient will also be subjected to an ultrasound examination of the knee for a maximum of 15 minutes. During this session a microbubble contrast, will be administered. For this session the burden for the subject will be minimal. For the patients who will undergo knee surgery, synovial tissue and synovial liquid will be collected during surgery. This is an additional procedure during surgery, but with a relatively low burden. During the MRI examination contrast fluid will be administered intravenously. Therefore an intravenous catheter is placed before the MRI scan. By this intravenous catheter a blood sample is taken for further tests. One vial will

be sufficient. The blood will be tested for example on biomarkers of inflammation.

Contacts

Public

Erasmus MC, Universitair Medisch Centrum Rotterdam

's Gravendijkwal 210

Rotterdam 3015 CE

NL

Scientific

Erasmus MC, Universitair Medisch Centrum Rotterdam

's Gravendijkwal 210

Rotterdam 3015 CE

NL

Trial sites

Listed location countries

Netherlands

Eligibility criteria

Age

Adults (18-64 years)

Elderly (65 years and older)

Inclusion criteria

- Age * 18 years;
- Clinical signs of knee OA: knee pain (numeric rating scale > 3);
- Radiographic signs of knee osteoarthritis (radiographic Kellgren and Lawrence grade *1);
- Signs of synovitis (i.e. joint effusion) on physical examination of the knee;

Exclusion criteria

- Subjects with a typical (e.g. metallic foreign bodies, etc.) contra-indication for an MRI exam;
- Renal insufficiency, checked with blood sample test (GFR < 60 ml/min/1,73 m²);
- Known allergy to contrast agents;
- Known pulmonary hypertension
- Woman who are pregnant or lactating

Study design

Design

Study type: Observational invasive

Masking: Open (masking not used)

Control: Uncontrolled

Primary purpose: Diagnostic

Recruitment

NL

Recruitment status: Recruiting

Start date (anticipated): 18-05-2017

Enrollment: 60

Type: Actual

Ethics review

Approved WMO

Date: 30-06-2016

Application type: First submission

Review commission: METC Erasmus MC, Universitair Medisch Centrum Rotterdam (Rotterdam)

Approved WMO

Date: 27-06-2017

Application type: Amendment

Review commission: METC Erasmus MC, Universitair Medisch Centrum Rotterdam (Rotterdam)

Approved WMO	
Date:	04-09-2017
Application type:	Amendment
Review commission:	METC Erasmus MC, Universitair Medisch Centrum Rotterdam (Rotterdam)
Approved WMO	
Date:	12-10-2017
Application type:	Amendment
Review commission:	METC Erasmus MC, Universitair Medisch Centrum Rotterdam (Rotterdam)

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register	ID
CCMO	NL57741.078.16