

Evaluation and testing of the microvascular function in asymptomatic athletes

Published: 10-02-2017

Last updated: 20-04-2024

The main objective of this study is to evaluate whether myocardial ischemia in the absence of obstructive CAD in athletes is associated with a reduced myocardial perfusion when compared to athletes with a normal exercise test. The secondary...

Ethical review	Not approved
Status	Will not start
Health condition type	Myocardial disorders
Study type	Observational invasive

Summary

ID

NL-OMON46277

Source

ToetsingOnline

Brief title

ESTIMATE Study

Condition

- Myocardial disorders

Synonym

Microvasculair function, the function of the smaller cardiac blood vessels

Research involving

Human

Sponsors and support

Primary sponsor: Maxima Medisch Centrum

Source(s) of monetary or material Support: Subsidie aangevraagd bij: Stichting Vrienden van het Hart Zuidoost Brabant. Sport Medisch Advies Centrum Veldhoven

Intervention

Keyword: Athletes, Microvascular function, PET-CT

Outcome measures

Primary outcome

Main study parameters:

- * Myocardial blood flow (MBF) per gram of myocardial tissue (ml/min/g) with ^{82}Rb PET-CT imaging.

- * Myocardial flow reserve (MBF at peak hyperaemia / MBF at rest) with ^{82}Rb PET-CT imaging.

Secondary outcome

Secondary study parameters:

- * The plasma levels of eNOS and ET-1 in athletes with a normal and abnormal exercise test result.

Study description

Background summary

Regular aerobic exercise has been shown to reduce the risk for fatal and non-fatal cardiac events and is therefore highly recommended both for healthy subjects and patients with cardiovascular disease. However, in selected cases, acute vigorous exercise may also be the trigger for a potential fatal cascade leading to myocardial infarction or sudden cardiac death (SCD). In order to reduce the incidence of SCD, the use of pre-participation screening is recommended for athletes to detect potential causes of SCD at an early stage. Exercise electrocardiography is particularly recommended to detect obstructive coronary artery disease (CAD). However, in many athletes (up to 95%) with an abnormal exercise test result, no obstructive CAD is found. The etiology and

prognostic implications of an abnormal exercise test in the absence of obstructive CAD among athletes remains unknown. Based on preliminary (animal) studies possible explanations may be an inadequate increase in myocardial capillary density in response to development of training-induced myocardial hypertrophy or coronary microvascular dysfunction due to remodeling of intramural coronary arteries. Both mechanisms will eventually result in an insufficient myocardial blood supply during exercise and may therefore serve as a trigger for a cascade leading to myocardial infarction or potentially lethal arrhythmias.

Study objective

The main objective of this study is to evaluate whether myocardial ischemia in the absence of obstructive CAD in athletes is associated with a reduced myocardial perfusion when compared to athletes with a normal exercise test. The secondary objectives are to evaluate different levels of the precursor endothelial nitric oxide synthase (eNOS) and endothelin-1 (ET-1) between the two groups.

Study design

A single-center observational case-control study among asymptomatic recreational and competitive athletes that underwent pre-participation screening at the department of Sports Medicine of Máxima Medical Center and/or visited the department of Cardiology of Máxima Medical Center. Athletes with an abnormal exercise test and abnormal myocardial perfusion scintigraphy (MPS) indicating myocardial ischemia within the last five years will be selected. When sufficient athletes with abnormal test results are included, athletes with normal test results will be matched for age, BMI and type of sport (1:1 ratio). All included athletes will undergo a PET-CT and blood will be collected.

Study burden and risks

All athletes will undergo blood tests at Máxima Medical Center. Blood samples with a total of 4 tubes (13 mL of blood) will be collected to evaluate routine laboratory investigations (C-reactive protein, leucocyte count, kidney function, glucose and lipid profile) and the plasma levels of eNOS and ET-1. The risks of a direct venipuncture are hematoma and bleeding at the site of puncture, in very exceptional cases thrombophlebitis.

All athletes will undergo ⁸²Rb PET / CCTA and will be exposed to radiation with a total effective radiation dose of approximately 5.8 * 7.1 mSv. Prior to the ⁸²Rb PET / CCTA, two intravenous lines will be placed which are both required (1 line for administration of ⁸²Rubidium and 1 line for administration of Adenosine). Risks of an intravenous line are minimal and comparable to a normal direct venipuncture (see above). Adverse effects that are described during

adenosine administration are bradycardia, premature atrial or ventricular beats, dyspnea, blushing, nausea and vomiting. All side effects disappear when administration is discontinued. Athletes are continuously monitored during administration and if serious side effects occur; administration will be discontinued immediately.

All participating athletes will be subjected to an effective radiation dose of $5.8 * 7.1$ mSv. When compared to the 2.6 mSv annual radiation exposure from natural sources in the Netherlands, this is a 2.2 to 2.7-fold increase in radiation exposure. A previous study showed that there is an increased non-fatal cancer risk of 0.01% per mSv of exposure to radiation, leading to an increased risk of non-fatal cancer of $0.058 * 0.071\%$ in the participating athletes. In order to avoid a high cumulative dose the use of radiation should be minimized. In this perspective, other non-invasive modalities to investigate the microcirculation were considered. Perfusion MRI is one of these modalities. The main advantage of this procedure is that MRI does not use ionizing radiation. However, PET is currently the golden standard for the assessment of microvascular function due to the highly accurate myocardial blood flow quantification. In contrast with ^{82}Rb PET, absolute measures of rest and stress flow with perfusion MRI did not correlated well. Also, robust fully-quantitative models have been developed only in experimental settings. Due to these limitations, visual assessment of the myocardial perfusion remains clinically the most used analysis tool when cardiac MRI is used. In the present study, the main study parameters are myocardial blood flow and flow reserve. As accurate quantitative measurements are needed, these parameters can only be obtained via PET- ^{82}Rb in the current clinical practice.

Coronary CT will be used to obtain anatomical imaging of the coronary arteries. This non-invasive modality can accurately detect and exclude the presence of epicardial CAD. As epicardial CAD may induce an impaired myocardial blood flow, it is necessary to rule out its presence. In this way, the calculated myocardial blood flow and flow reserve via ^{82}Rb PET / CCTA will solely reflect the microvascular function. The index athletes underwent coronary angiography (CAG) as part of the diagnostic procedures. However, this CAG is performed months to years before recruitment for the present study. Therefore, as CAD at the time of study enrollment cannot be ruled out with certainty, the coronary anatomy will be evaluated also at the time of study enrollment. For this purpose, ^{82}Rb PET will be combined with a CCTA scan, which imposes an additional radiation dose of $3.4 * 4.1$ mSv which is lower than the radiation dose of CAG ($5 * 7$ mSv).

Although there is no major benefit for individual study participants, the results of this study may lead to a better understanding of microvascular coronary (dys)function in asymptomatic athletes on a population level, and consequently, to an increased insight whether athletes with positive exercise test result are prone to suffer cardiac (lethal) events and should be advised to refrain from competitive sports. In case of abnormal findings (e.g. coronary

artery disease), the subject will be referred to their own general practitioner with a recommendation for referral to a cardiologist. If the subject is currently under treatment of a cardiologist, a direct referral will be initiated and the subject will be treated according to the current standards.

Contacts

Public

Maxima Medisch Centrum

De Run 4600
Veldhoven 5504 DB
NL

Scientific

Maxima Medisch Centrum

De Run 4600
Veldhoven 5504 DB
NL

Trial sites

Listed location countries

Netherlands

Eligibility criteria

Age

Adults (18-64 years)
Elderly (65 years and older)

Inclusion criteria

Asymptomatic athletes with normal exercise electrocardiography results and asymptomatic athletes with concordant abnormal exercise electrocardiography results and abnormal myocardial perfusion scintigraphy results but without the presence of epicardial coronary artery disease.;An athlete is defined as a sportsman who is engaged in sports for at least 2.5 hours a week for a period of minimally 30 weeks per year or in two or more sports with a minimum of 1.5 hours per week within one type of sports for at least 20 weeks per year

Exclusion criteria

Symptomatic athletes ((Exercise induced) chest pain, palpitations, dizziness, light headiness or syncope), athletes with epicardial coronary artery disease, athletes with previous myocardial infarction

Study design

Design

Study type:	Observational invasive
Intervention model:	Other
Allocation:	Non-randomized controlled trial
Masking:	Open (masking not used)
Control:	Active
Primary purpose:	Basic science

Recruitment

NL	
Recruitment status:	Will not start
Enrollment:	50
Type:	Anticipated

Ethics review

Not approved	
Date:	10-02-2017
Application type:	First submission
Review commission:	METC Maxima Medisch Centrum (Veldhoven)

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register	ID
CCMO	NL55136.015.16