Determining Exercise Tolerance using an EMG Cycle Test in patients with a neuromuscular disorder

Published: 21-03-2019 Last updated: 09-04-2024

Gaining an objective insight in muscle fatigue during exercise for patients with NMD by1. comparing the timing of change in sEMG amplitude and sEMG median frequency, to gas exchange variables, heart rate, work load, muscle strength, muscular...

Ethical review	Approved WMO	
Status	Recruitment stopped	
Health condition type	Musculoskeletal and connective tissue disorders congenital	
Study type	Observational invasive	

Summary

ID

NL-OMON48098

Source ToetsingOnline

Brief title DETECT-NMD

Condition

Musculoskeletal and connective tissue disorders congenital

Synonym

muscle diseases, neuromuscular disorders

Research involving Human

Sponsors and support

Primary sponsor: Radboud Universitair Medisch Centrum Source(s) of monetary or material Support: stichting afdeling Revalidatie Radboudumc

1 - Determining Exercise Tolerance using an EMG Cycle Test in patients with a neurom ... 31-05-2025

Intervention

Keyword: EMG, Exercise tolerance, Neuromuscular disorders, Outcome measure

Outcome measures

Primary outcome

sEMG amplitude

sEMG median frequency

Secondary outcome

3D movement analysis

spirometry (VO2 max, heart rate, anaerobic threshold, workolad, serum lactate

levels, Borg scale)

Muscle strength

Study description

Background summary

On average more than 60% of patients with a neuromuscular disorder (NMD)experiences severe fatigue. Increasing the level of physical inactivity by aerobic exercises or cognitive behavioural therapy leads to a lower level of fatigue and a deceleration of the increase of fatty infiltration of upper leg muscles in the majority of these patients. However, the effect of both therapies on exercise tolerance is unknown. Current outcome measures for exercise tolerance focus on the cardiovascular system by the use of ergospirometry. For patients with a NMD , muscle fatigue generally is the limiting

factor during exercise. One possible non-invasive alternative to objectively measure muscle fatigue on the (neuro) muscular level could be the use of surface electromyography (sEMG).

Study objective

Gaining an objective insight in muscle fatigue during exercise for patients with NMD by

1. comparing the timing of change in sEMG amplitude and sEMG median frequency, to gas exchange variables, heart rate, work load, muscle strength, muscular oxidative capacity and blood lactate levels during a cycle test with a constant and incremental workload en strength measurements of arms and legs

2. comparing the relative timing of change in sEMG amplitude and sEMG median frequency in patients with NMD with

the relative timing of change in sEMG amplitude and sEMG median frequency in healthy subjects

Study design

Explorative cross-sectional proof of concept pilot study

Study burden and risks

The burden and risks associated with participation are neglectable. The task that need to be performed is safely performed by healthy subjects and NMD patients before. CE marked equipment is used.

Contacts

Public

Radboud Universitair Medisch Centrum

Reinier postlaan 2 Nijmegen 6525 GC NL **Scientific** Radboud Universitair Medisch Centrum

Reinier postlaan 2 Nijmegen 6525 GC NL

Trial sites

Listed location countries

Netherlands

3 - Determining Exercise Tolerance using an EMG Cycle Test in patients with a neurom ... 31-05-2025

Eligibility criteria

Age

Adults (18-64 years) Elderly (65 years and older)

Inclusion criteria

- aged 18 years and older
- being able to read and understand the Dutch language
- mentally competent
- being able to cycle at least 8 minutes on a cycle leg and/ or arm ergometer

Exclusion criteria

- pregnancy
- cognitive impairment
- disabling co-morbidity interfering with the cycle test
- contraindications to Exercise Stress Testing according to the guidelines of the American Heart Association

Study design

Design

Study type:	Observational invasive
Intervention model:	Other
Allocation:	Non-randomized controlled trial
Masking:	Open (masking not used)
Control:	Active
Primary purpose:	Diagnostic

Recruitment

NII

NL .	
Recruitment status:	Recruitment stopped
Start date (anticipated):	21-03-2019

4 - Determining Exercise Tolerance using an EMG Cycle Test in patients with a neurom ... 31-05-2025

Enrollment:	90
Туре:	Actual

Ethics review

Approved WMO	
Date:	21-03-2019
Application type:	First submission
Review commission:	CMO regio Arnhem-Nijmegen (Nijmegen)

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register ССМО

ID NL68723.091.19