

# Sentinel lymph node detection in early-stage oral cavity squamous cell carcinoma using computed tomographic lymphography

Published: 20-04-2020

Last updated: 08-04-2024

The primary objective of this study is to assess the diagnostic accuracy, in terms of sensitivity and negative predictive value, of preoperative CT lymphography as compared with conventional lymphoscintigraphy for SLN detection. Secondly, we aim to...

<b>Ethical review</b>	Approved WMO
<b>Status</b>	Recruitment stopped
<b>Health condition type</b>	Head and neck therapeutic procedures
<b>Study type</b>	Observational invasive

## Summary

### ID

NL-OMON49703

### Source

ToetsingOnline

### Brief title

SELECT

### Condition

- Head and neck therapeutic procedures

### Synonym

Mouth cancer, occult lymph node metastasis

### Research involving

Human

### Sponsors and support

**Primary sponsor:** Universitair Medisch Centrum Utrecht

**Source(s) of monetary or material Support:** Ministerie van OC&W

## **Intervention**

**Keyword:** CT lymphography, Oral cavity carcinoma, Sentinel lymph node biopsy

## **Outcome measures**

### **Primary outcome**

The sensitivity and negative predictive value of CT lymphography as compared with conventional lymphoscintigraphy for SLNB. In addition, the sensitivity and negative predictive value for preoperative CT lymphography combined with conventional preoperative lymphoscintigraphy.

### **Secondary outcome**

- To compare the number of CT lymphographic detected SLNs with those detected by means of 99mTc-nanocolloid lymphoscintigraphy on a per-subject basis.
- To compare the number of CT lymphographic detected higher-echelon nodes with those detected by means of 99mTc-nanocolloid lymphoscintigraphy on a per-subject basis.
- To compare histopathologic assessment (presence or absence of metastasis) of the excised lymph node(s) detected by conventional preoperative 99mTc-nanocolloid lymphoscintigraphy and intraoperative gammaprobe localization, with the SLNs identified by means of preoperative CT lymphography.
- Observing contralateral drainage patterns in lateralized tumors and compare these patterns between CT lymphography and conventional 99mTc-nanocolloid lymphoscintigraphy.
- To assess pairwise inter-observer agreements between CT lymphography and

conventional  $^{99m}\text{Tc}$ -nanocolloid lymphoscintigraphy regarding preoperative SLN detection.

## Study description

### Background summary

Cervical lymph node metastasis is the single most important prognostic factor in head and neck squamous cell carcinoma (HNSCC), including oral cavity squamous cell carcinoma (OSCC), and accurate detection of cervical lymph node metastases is critical for surgical and adjuvant therapy planning and prognosis prediction. The sentinel lymph node (SLN) procedure is a diagnostic staging procedure that is applied in a variety of tumour types, including HNSCC. The SLN is the first draining lymph node from the tumor, which is most likely to harbour metastases. The histopathological status of the SLN should reflect the histopathological status of the rest of the nodal basin, and additional treatment of the nodal basin (e.g., surgery) should be performed in case of metastatic involvement of the SLN. Detecting SLNs close to tumor sites is hampered, since the injection site of the radiotracer, around the primary tumor, produces a large hotspot on lymphoscintigraphy possibly hiding SLNs in close proximity of the primary tumor (\*shine through\* effect). SLN imaging using CT techniques might be more sensitive for detection of SNLs.

### Study objective

The primary objective of this study is to assess the diagnostic accuracy, in terms of sensitivity and negative predictive value, of preoperative CT lymphography as compared with conventional lymphoscintigraphy for SLN detection. Secondly, we aim to assess the diagnostic accuracy of preoperative CT lymphography combined with conventional preoperative lymphoscintigraphy for SLN detection.

### Study design

1. A pilot study to optimize the CT lymphography imaging protocol and build experience with the outcomes of CT lymphography (10 patients).
2. A prospective cohort study and a within-patient evaluation of CT lymphography for identification of SLNs as compared to conventional  $^{99m}\text{Tc}$ -nanocolloid lymphoscintigraphy in patients with early-stage OSCC (84 patients).

### Study burden and risks

Patients will undergo an additional CT lymphography after peritumoral

injections with lipiodol. The information obtained by CT lymphography may provide helpful anatomical information of the SLN(s) for the surgeon, which may help in harvesting the SLN(s), especially in cases where the SLN(s) are located close to the injection site of the primary tumor. The extra administration of 0.5 mL lipiodol, followed by CT imaging, will not result in a significant additional (radiation) burden for the patient.

## Contacts

### Public

Universitair Medisch Centrum Utrecht

Heidelberglaan 100  
Utrecht 3584 CX  
NL

### Scientific

Universitair Medisch Centrum Utrecht

Heidelberglaan 100  
Utrecht 3584 CX  
NL

## Trial sites

### Listed location countries

Netherlands

## Eligibility criteria

### Age

Adults (18-64 years)

Elderly (65 years and older)

### Inclusion criteria

In order to be eligible to participate in this study, a subject must meet all of the following criteria:

1. The patient has provided written informed consent authorization before participating in the study.

2. The patient has a diagnosis of primary oral squamous cell carcinoma that is anatomically located in: mucosal lip, buccal mucosa, lower alveolar ridge, upper alveolar ridge, retromolar gingival (retromolar trigone), floor-of-the-mouth, hard palate or oral (mobile) tongue, and is stage T1-T2 and T3 (only when T3 is assessed based on tumour dimensions of >2 cm and ≤4 cm with DOI >10 mm), N0, M0.
3. Clinical nodal staging (N0) has been confirmed by negative results from ultrasound guided fine needle aspiration cytology.
4. The patient is a candidate for transoral excision and sentinel lymph node biopsy.
5. The patient has given informed consent for a surgical procedure regarding his/her oncological treatment.
6. Patients with prior malignancy of the head and neck area are allowed, provided the patient meets both of the following criteria:
  - Underwent potentially curative therapy for all prior head and neck malignancies and is deemed low risk for recurrence; and
  - No head and neck malignancy for the past five years and no evidence of recurrence.
7. The patient is >18 years of age at the time of consent.
8. The patient has an ECOG status of Grade 0 - 2.

## Exclusion criteria

1. The patient has a diagnosis of squamous cell carcinoma of the head and neck in the following anatomical areas: non-mobile base of the tongue, oropharynx, nasopharynx, hypopharynx, and larynx.
2. Patient is incapacitated.
3. Patient is pregnant or lactating.
4. Previous allergic reaction after administration of a CT-contrast-agent for contrast enhanced CT-imaging.
5. Patient is known with manifest hyperthyroidism.
6. The patient has clinical or radiological evidence of metastatic cancer to the regional lymph nodes.
7. The patient has a history of neck dissection, or gross injury to the neck that would preclude reasonable surgical dissection for this trial, or radiotherapy to the neck.
8. The patient is actively receiving systemic cytotoxic chemotherapy.
9. Patient is on immunosuppressive, anti-monocyte, or immunomodulatory therapy.
10. Participation will result in unacceptable delay regarding oncological treatment.

## Study design

### Design

**Study type:** Observational invasive

Masking: Open (masking not used)

Control: Uncontrolled

Primary purpose: Diagnostic

### Recruitment

NL

Recruitment status: Recruitment stopped

Start date (anticipated): 09-11-2020

Enrollment: 94

Type: Actual

## Ethics review

Approved WMO

Date: 20-04-2020

Application type: First submission

Review commission: METC NedMec

## Study registrations

### Followed up by the following (possibly more current) registration

No registrations found.

### Other (possibly less up-to-date) registrations in this register

No registrations found.

## In other registers

### Register

CCMO

### ID

NL72330.041.20