

In-vivo multispectral optoacoustic imaging of thyroid nodules

Published: 27-11-2019

Last updated: 19-08-2024

We aim to determine if in vivo MSOT can distinguish benign from malignant thyroid nodules by correlating and validating optoacoustic signals in vivo with pathology results.

Ethical review	Approved WMO
Status	Recruitment stopped
Health condition type	Endocrine neoplasms malignant and unspecified
Study type	Interventional

Summary

ID

NL-OMON49819

Source

ToetsingOnline

Brief title

THYNODE

Condition

- Endocrine neoplasms malignant and unspecified

Synonym

thyroid nodules, thyroid tumor

Research involving

Human

Sponsors and support

Primary sponsor: Universitair Medisch Centrum Groningen

Source(s) of monetary or material Support: Ministerie van OC&W

Intervention

Keyword: MSOT, Multispectral, Optoacoustic, Thyroid nodules

Outcome measures

Primary outcome

The main study endpoint is quantification of the optoacoustic signal (HbO₂, HbR, HbT, sO₂, fat, water and collagen) observed by multispectral optoacoustic imaging using the MSOT Acuity Echo in vivo in patients with thyroid nodules, also in relation to the definitive pathology results (benign vs malignant).

Secondary outcome

- To explore a potential correlation between endogenous optoacoustic signals in vivo with cytology (the standard-of-care of Bethesda scoring system derived from FNA)
- To explore a potential correlation between endogenous optoacoustic signals in vivo with TIRADS ultrasonography scoring system (the scoring system used by radiologists when evaluating the thyroid nodules with ultrasonography).

Study description

Background summary

Thyroid nodules are common in clinical practice. Head and neck ultrasound is recommended as a routine examination for all patients with thyroid lesions. The Thyroid Imaging Reporting And Documentation System (TIRADS) criteria helps to estimate the risk of malignancy based on ultrasound patterns and nodule sizes guiding the performance of fine-needle-aspiration (FNA). Approximately 20% of FNA results cannot be specified whether being benign or malignant tissue. A definitive diagnosis can only be made from histopathology after diagnostic (hemi)thyroidectomy. However, (hemi)thyroidectomy has disadvantages as it leads to over-treatment and has a risk of postoperative morbidity (e.g. hypothyroidism and laryngeal nerve injury). Furthermore, (hemi)thyroidectomy is known to be associated with poor quality of life. Clearly, there is an unmet need for additional diagnostic tools in order to identify malignant thyroid nodules and thereby support the decision making for treatment of the thyroid. Multispectral optoacoustic tomography (MSOT) is a novel non-invasive imaging

method that enables visualization of endogenous chromophores and exogenous contrast agents using the generation of ultrasound waves due to light absorption. Recently, this system has been used for non-invasive determination of thyroid nodules. Results show that multispectral optoacoustic imaging of thyroid nodules may distinguish benign from malignant nodules. However, a larger cohort is necessary to confirm this finding.

Study objective

We aim to determine if in vivo MSOT can distinguish benign from malignant thyroid nodules by correlating and validating optoacoustic signals in vivo with pathology results.

Study design

The current study is a non-randomized, non-blinded, single center, prospective, cross-sectional, proof of concept study.

Intervention

Included patients will undergo MSOT imaging of the thyroid nodules.

Study burden and risks

A potential risk of MSOT is cornea damage when looking straight into the laser. As everyone present in the MSOT room during imaging is obligated to wear laser safety goggles, this risk is covered. In addition, there is an interlock system in the room so that no one else can enter the room when the imaging system is activated.

The risk associated with the use of MSOT seems minor and although patients will not directly benefit from this study, results of this study will provide valuable information for our understanding of the diagnostic value of MSOT in distinguishing benign from malignant thyroid nodules.

Contacts

Public

Universitair Medisch Centrum Groningen

Hanzeplein 1
Groningen 9700 RB
NL

Scientific

Universitair Medisch Centrum Groningen

Hanzeplein 1
Groningen 9700 RB
NL

Trial sites

Listed location countries

Netherlands

Eligibility criteria

Age

Adults (18-64 years)

Elderly (65 years and older)

Inclusion criteria

1. Patients with thyroid nodules who underwent or will undergo an ultrasonography (with TIRADS score) with FNA if indicated and will be scheduled for a (hemi)thyroidectomy if indicated;
2. Age ≥ 18 years;
3. Written informed consent.

Exclusion criteria

1. Medical or psychiatric conditions that compromise the patients' ability to give informed consent;
2. Previous surgery in head and neck area on the ipsilateral side of the index nodule
3. Previous radiotherapy in head and neck area
4. Pregnancy

Study design

Design

Study type: Interventional

Masking: Open (masking not used)

Control: Uncontrolled

Primary purpose: Diagnostic

Recruitment

NL

Recruitment status: Recruitment stopped

Start date (anticipated): 18-09-2020

Enrollment: 30

Type: Actual

Medical products/devices used

Generic name: Multispectral optoacoustic tomography

Registration: No

Ethics review

Approved WMO

Date: 27-11-2019

Application type: First submission

Review commission: METC Universitair Medisch Centrum Groningen (Groningen)

Approved WMO

Date: 25-11-2020

Application type: Amendment

Review commission: METC Universitair Medisch Centrum Groningen (Groningen)

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register	ID
CCMO	NL69993.042.19