Articulation and coordination of speech after treatment for oral cancer

Published: 25-05-2021 Last updated: 08-04-2024

To investigate the timing and coordination of speech articulation and its relation to speech disorders of people who have been treated for oral cancer with a major resection. This information will lead to a better understanding of articulation and...

Ethical review	Approved WMO
Status	Recruitment stopped
Health condition type	Head and neck therapeutic procedures
Study type	Observational invasive

Summary

ID

NL-OMON51026

Source ToetsingOnline

Brief title Speech difficulties after oral cancer treatment

Condition

· Head and neck therapeutic procedures

Synonym dysarthria, speech pathology

Research involving Human

Sponsors and support

Primary sponsor: Universitair Medisch Centrum Groningen Source(s) of monetary or material Support: Ministerie van OC&W

Intervention

Keyword: Articulation, dysarthria, Oral cancer, Speech

Outcome measures

Primary outcome

- Displacement of the tongue tip, the tongue body, the lips and the jaw at

consonant and vowel targets

- Velocity of tongue, of the tongue tip, the tongue body, the lips, and the jaw
- at key landmarks of speech gestures.
- Duration of speech gestures produced with the tongue tip, the tongue body,

the lips and the jaw.

- Speech rate (syllables/min and words/min)

Secondary outcome

- Acoustic measures of speech
- Models to link articulation and speech, both speech synthesis from EMA, and

EMA from speech (machine learning)

- Changes in the variability and complexity of articulators* movement under

masked auditory feedback compared to normal feedback

Study description

Background summary

People treated for oral cancer with a major resection procedure often develop problems with swallowing and articulation. The deviant articulation affects speech, complicates communication, and has a negative effect on the quality of life of patients. Currently, the available information on the effects of oral cancer treatment on speech articulation is rather scarce, especially with respect to the changes and adaptations in articulatory processes in these speakers. The proposed research will shed more light on this issue and will lead ultimately to a better understanding of the anatomical and physiological basis of the disordered speech in oral cancer patients. In turn, this information will be-come available to clinicians and speech therapists who will be able to minimise negative effects of treatment and improve therapies after treatment.

Study objective

To investigate the timing and coordination of speech articulation and its relation to speech disorders of people who have been treated for oral cancer with a major resec-tion. This information will lead to a better understanding of articulation and articulation related speech problems after oral cancer and will ultimately aid speech therapy and improve the information and consultation of these patients.

Study design

Subjects will perform several speech tasks. Some of the speech tasks will be carried out while the participant is wearing earphones through which pink noise is distributed (at a comfortable volume). During these tasks, articulatory trajectories will be recorded using electromagnetic articulography (EMA). The data from the oral cancer patients will be compared to EMA and speech data from non-speech disturbed controls.

Study burden and risks

No known risks or benefits are associated with participating in this study.

Contacts

Public Universitair Medisch Centrum Groningen

Hanzeplein 1 Groningen 9713GZ NL **Scientific** Universitair Medisch Centrum Groningen

Hanzeplein 1 Groningen 9713GZ NL

Trial sites

Listed location countries

Netherlands

Eligibility criteria

Age

Adults (18-64 years) Elderly (65 years and older)

Inclusion criteria

- At least 18 years of age and able to make an informed consent
- Native speaker of Dutch

• Diagnosed with oral tumors and having had a major surgical resection of a TNM stage T3 or T4, where parts of the tongue or jaw have been removed and perhaps replaced by other tissue at least 12 months before (patient subject) or a volunteer without disturbed speech.

Exclusion criteria

- Recurrence of disease
- History of neurological or psychological disorders
- Self-reported signs of depression
- Stuttering or other pre-existing speech and language problems
- Anatomy that prevents attaching sensors to the tongue (i.e., trismus or tongue immobility)

• Problems with sight or hearing that impede reading or understanding instructions. When glasses or a hearing aid resolve these problems, there is no impediment to participation.

• Non removable metal on, in or close to the head (e.g., piercings, dental braces, medical devices such as deep brain stimulation electrodes) or medical devices (e.g., a pacemaker) incompatible to electromagnetic fields.

Study design

Design

Study type:	Observational invasive
Intervention model:	Other
Allocation:	Non-randomized controlled trial
Masking:	Open (masking not used)

Primary purpose: Diagnostic

Recruitment

NI

Recruitment status:	Recruitment stopped
Start date (anticipated):	28-09-2021
Enrollment:	40
Туре:	Actual

Ethics review

Approved WMO	
Date:	25-05-2021
Application type:	First submission
Review commission:	METC Universitair Medisch Centrum Groningen (Groningen)

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register CCMO **ID** NL76137.042.20