

# Assessment of Treatment with Laparoscopic fenestration or Aspiration Sclerotherapy for large symptomatic hepatic cysts (ATLAS): a randomized clinical trial.

Published: 14-06-2021

Last updated: 15-05-2024

The main objective is to compare laparoscopic fenestration and aspiration sclerotherapy in patients with large symptomatic hepatic cysts on patient-reported outcomes. This information can be used to assess cost-effectiveness in both treatments.

<b>Ethical review</b>	Approved WMO
<b>Status</b>	Will not start
<b>Health condition type</b>	Hepatobiliary disorders congenital
<b>Study type</b>	Interventional

## Summary

### ID

NL-OMON51055

### Source

ToetsingOnline

### Brief title

ATLAS-trial

### Condition

- Hepatobiliary disorders congenital
- Hepatic and hepatobiliary disorders
- Hepatobiliary therapeutic procedures

### Synonym

hepatic cysts

### Research involving

Human

## Sponsors and support

**Primary sponsor:** Universitair Medisch Centrum

**Source(s) of monetary or material Support:** Ministerie van OC&W

## Intervention

**Keyword:** aspiration sclerotherapy, Hepatic cyst, laparoscopic fenestration

## Outcome measures

### Primary outcome

Main study parameters: The main study parameter is the PLD-Q score at 4 weeks after treatment.

### Secondary outcome

Secondary parameters are among others: PLD-Q score at baseline, 6 months and 12 months; liver volume (CT) at baseline and 4 weeks; cyst volume (ultrasound) at baseline, 4 weeks, 6 months and 12 months; complications according to Clavien-Dindo; admission duration, recurrence and re-intervention rates.

## Study description

### Background summary

Patients with large hepatic cysts(>5cm) may develop symptoms due to distention of Glisson's capsule and/or compression on other abdominal organs. Frequently reported symptoms include abdominal pain, early satiety, nausea, and dyspnea. These symptoms can be captured in the disease-specific Polycystic Liver Disease Questionnaire (PLD-Q), a validated instrument. The treatment of symptomatic liver cysts is aimed to improve symptoms and quality of life by reducing cyst volume. There are two procedures available to treat symptomatic liver cysts: percutaneous aspiration sclerotherapy and laparoscopic fenestration.

In aspiration sclerotherapy, fluid is evacuated from the liver cyst and subsequently the cyst lining is exposed to a sclerosing agent for a limited

period of time. Sclerotherapy causes temporary recurrence of cyst fluid after drainage, but subsequently results in a steady decrease of cyst volume in the majority of patients.

In laparoscopic fenestration the liver is exposed through laparoscopic surgery. In this procedure the cyst is punctured and drained followed by resection of extra-hepatic cyst wall.

The safety and efficacy of aspiration sclerotherapy and laparoscopic fenestration have been explored in two recent systematic reviews. No evident conclusion could be drawn because of the retrospective study design in the vast majority of the studies and the heterogeneity among these. A randomized controlled trial is warranted to identify the possible differences in safety and efficacy in aspiration sclerotherapy and laparoscopic fenestration.

Hypothesis: We expect patients treated with laparoscopic fenestration to have better clinical outcome; i.e. a lower PLD-Q score, compared to aspiration sclerotherapy, when measured 4 weeks after the procedure. We expect this difference to become smaller over time (after 6 and 12 months), with loss of statistical significance.

## **Study objective**

The main objective is to compare laparoscopic fenestration and aspiration sclerotherapy in patients with large symptomatic hepatic cysts on patient-reported outcomes. This information can be used to assess cost-effectiveness in both treatments.

## **Study design**

A single-center, prospective, randomized clinical superiority trial in which patients will be randomized 1:1 to one of the treatment arms. Patients will be followed for 1 year.

## **Intervention**

Intervention: Patients will be randomly allocated to either aspiration sclerotherapy or laparoscopic fenestration. Both procedures are performed according to the standard Radboudumc protocols. Aspiration sclerotherapy consists of ultrasound-guided, percutaneous drainage of the cyst with subsequent sclerosation with ethanol. Laparoscopic fenestration consists of standard abdominal laparoscopy in which the large cyst(s) are drained and deroofed.

## **Study burden and risks**

All symptomatic liver cysts patients that are included have an indication for treatment by aspiration sclerotherapy or laparoscopic fenestration. The described follow-up is according to the regular liver cyst treatment local protocol, including site visits, questionnaires and ultrasound. For patients allocated to the aspiration sclerotherapy arm of the study, one additional CT-scan is made at 4 weeks, which would not be standard protocol. In case of an inconclusive ultrasound at 6 months, a CT-scan (without contrast, low-dose) will be performed to measure cyst volume.

In summary, besides random allocation and extra follow-up CT-scan(s), the study forms no extra burden or associated risks. Associated risks in aspiration sclerotherapy are temporary and mild, e.g. local pain. Associated risks in laparoscopic fenestration are bile leakage, ascites, pleural effusion or infections. The number of questionnaire-sets that need to be filled in will be 4-5 (dependent on intermittent recurrence). The number of CT-scans that will be made during the study is 1, 2 or 3, depending on available imaging before screening and need for a CTs.

## Contacts

### **Public**

Selecteer

Geert Grooteplein Zuid 10  
Nijmegen 6525 GA  
NL

### **Scientific**

Selecteer

Geert Grooteplein Zuid 10  
Nijmegen 6525 GA  
NL

## Trial sites

### Listed location countries

Netherlands

## Eligibility criteria

## Age

Adults (18-64 years)

Elderly (65 years and older)

## Inclusion criteria

- Age  $\geq 18$  years
- Hepatic cyst characteristics:
  - Large ( $>5$  cm),
  - Symptomatic (PLD-Q score  $\geq 20$ ),
  - Non-parasitic on imaging (US/CT/MRI)
  - Non-neoplastic on imaging (US/CT/MRI)
- Providing informed consent

## Exclusion criteria

- Clinical suspicion of a complicated hepatic cyst (cyst rupture or active cyst infection)
- Cyst is not laparoscopically accessible for surgery
- Cyst is not percutaneously (ultrasound-guided) accessible for aspiration
- More than 20 cysts of  $>1.5$  cm
- Age above 70 years
- ASA IV
- ECOG score  $>1$
- Aspiration sclerotherapy or laparoscopic fenestration of hepatic cysts was performed in the last 6 months.
- Severe renal impairment (eGFR  $< 30$  ml/min/1,73 m<sup>2</sup>)
- Coagulopathy (spontaneous INR  $>2$  or platelet count  $< 80 \times 10^9/l$ )
- Radiologic contrast allergy
- Pregnancy
- Any current or prior medical condition that may interfere with the conduct of the study or the evaluation of its results in the opinion of the investigator (e.g. inability to fill out questionnaires).

## Study design

### Design

Study phase: 3

Study type: Interventional

Intervention model:	Parallel
Allocation:	Randomized controlled trial
Masking:	Open (masking not used)
Control:	Active
Primary purpose:	Treatment

## Recruitment

NL	
Recruitment status:	Will not start
Enrollment:	70
Type:	Anticipated

## Ethics review

Approved WMO	
Date:	14-06-2021
Application type:	First submission
Review commission:	CMO regio Arnhem-Nijmegen (Nijmegen)
Approved WMO	
Date:	18-11-2021
Application type:	Amendment
Review commission:	CMO regio Arnhem-Nijmegen (Nijmegen)

## Study registrations

### Followed up by the following (possibly more current) registration

No registrations found.

### Other (possibly less up-to-date) registrations in this register

ID: 20962  
Source: Nationaal Trial Register  
Title:

## In other registers

Register	ID
CCMO	NL75717.091.20
OMON	NL-OMON20962