

# NETosis in burn wound patients

Published: 14-04-2023

Last updated: 16-11-2024

In this clinical pilot study we want to gather data in order to calculate a power for correct subject group to answer the following research question: to explore the relation between NETs/cfDNA in the blood and NETosis in burn wounds and their...

<b>Ethical review</b>	Approved WMO
<b>Status</b>	Recruiting
<b>Health condition type</b>	Other condition
<b>Study type</b>	Observational invasive

## Summary

### ID

NL-OMON54153

### Source

ToetsingOnline

### Brief title

NETosis in burns/NIB

### Condition

- Other condition
- Embolism and thrombosis

### Synonym

NETosis and thrombosis formation

### Health condition

Brandwonden

### Research involving

Human

### Sponsors and support

**Primary sponsor:** Vrije Universiteit Medisch Centrum

**Source(s) of monetary or material Support:** Nederlandse brandwonden stichting;Health

## Intervention

**Keyword:** Burn wounds, NETosis, Thrombosis, Wound expansion.

## Outcome measures

### Primary outcome

Burn wound severity will be determined at day of admission (day 0/1), day 2-5 post-burn and at day of surgery (if applicable) or day 10 - 20 post-burn by clinical evaluation, Laser Doppler Imaging (LDI) (not day 10 - 20), and (immuno)histochemical analyses in burn skin biopsies. In biopsies taken at the same time points the occurrence of intra-microvascular thrombi and NETosis will be quantified. In addition, in blood samples collected simultaneously with clinical routine lab up to day 30 post-burn, the levels of cfDNA/NETs, DNase1 and endogenous pdC1inh will be quantified. MBL serum levels will be examined in relation to bacterial presence and wound healing.

From healthy volunteers only blood samples will be collected for the quantification of cfDNA/NETs, DNase1 and endogenous pdC1inh levels.

### Secondary outcome

Correlations/associations between wound deepening, thrombosis/NETosis, levels of cfDNA/NETs, DNase1 and endogenous pdC1inh will be investigated. In addition, associations of thrombosis/NETosis with burn wound parameters and characteristics of the patients collected from the medical file e.g. wound size (% burned of TBSA), wound closure time, co-morbidities, inhalation trauma, graft take, length of stay, number of surgical procedures will also be

investigated.

## Study description

### Background summary

Upon burn injury a massive inflammatory reaction is induced, which is initially essential to combat invading microorganisms, debride the wound from damaged tissue and regulate the wound healing. However, in burn wound patients this massive inflammatory response seems to be \*uncontrolled\* and persist for up to months after the initial trauma, which not only negatively affects the local healing process of the burn wound, but additionally exerts systemic effects may result in e.g. secondary (organ) injury and burn wound extension. In addition, prevention of microbial contamination and infection is vital for burn wound care. Bacterial presence can namely result in further wound healing problems.

Part 1. Inflammation and burn wound extension: Neutrophil extracellular traps in expansion of necrosis of the burn wound In burn wound patients, the expansion of partial thickness burn wounds to deeper or full thickness burn wounds in the first 48 hours is a common phenomenon. This post-burn wound deepening leads to increased tissue loss, delayed healing, more hypertrophic scarring and contractures, an increased need for surgical excisions and grafting and an increased chance of burn wound infections and death. Despite the use of e.g. cerium nitrate and anticoagulants in order to prevent thrombosis, so far no effective treatment exists to prevent post-burn wound deepening. We recently found extensive microvascular thrombosis in and around burn wounds, in both animals and human tissue, which persisted for up to weeks after the initial burn injury. This coincided with neutrophil extracellular traps (NETs) formation (i.e. NETosis), the process whereby neutrophils eject net-like structures of their DNA. NETosis has been shown in general to be a major initiator of blood coagulation and microvascular damage and may therefore be a promising therapeutic target in order to prevent microvascular thrombosis and wound deepening in burn wound patients.

Part 2. Immunological factors of infection

In burn wounds, *Staphylococcus aureus* and *Pseudomonas aeruginosa* are the most frequently isolated microbial species. Deficiency for mannose-binding lectin (MBL) may predispose patients to colonization and infection. MBL is a broad-spectrum pattern recognition molecule that plays a role in innate immunity and in the inflammatory response after skin injury. A low serum level of MBL together with other comorbid factors may predispose the host to increased susceptibility to colonization and infection.

### Study objective

In this clinical pilot study we want to gather data in order to calculate a

power for correct subject group to answer the following research question: to explore the relation between NETs/cfDNA in the blood and NETosis in burn wounds and their prognostic potential with regards to clinical outcome. In addition, we will investigate whether there are links between the concentration of MBL in blood, the presence of *P. aeruginosa* and/or *S. aureus* in burns and wound healing problems.

## **Study design**

Prospective observational cohort pilot study.

## **Study burden and risks**

The burden of this study is burned skin biopsy collection, LDI and clinical assessments, daily/weekly blood withdrawals (combined with clinical routine sampling as much as possible).

The collection of skin biopsies will take place at day of admission (day 0/1), at day 2 - 5 post-burn and at day of surgery (if applicable) or at day 10 - 20 post-burn.

Clinical parameters e.g. pain and wound healing parameters will be assessed during routine clinical assessments.

The burden of the study for healthy volunteers is limited to two blood withdrawals via venepunctures.

## **Contacts**

### **Public**

Vrije Universiteit Medisch Centrum

De Boelelaan 1108  
Amsterdam 1081 HZ  
NL

### **Scientific**

Vrije Universiteit Medisch Centrum

De Boelelaan 1108  
Amsterdam 1081 HZ  
NL

## **Trial sites**

## Listed location countries

Netherlands

## Eligibility criteria

### Age

Adults (18-64 years)

Elderly (65 years and older)

## Inclusion criteria

Patients:

- $\geq 18$  years
- Acute burns
- Competent and temporarily incompetent patients
- Informed Consent by patient or the legal representative
- Admitted to the Burn Center of the Red Cross Hospital in Beverwijk, who are expected to be admitted at least until the second measurement moment
- No acute psychiatric disorders
- Sufficient Dutch proficiency

Healthy volunteers:

- $\geq 18$  years
- Competent
- Informed Consent
- Sufficient Dutch proficiency

## Exclusion criteria

Patients:

- Known underlying diseases: cancer, immune deficiency, hypertension, complement, thrombosis disorder.

Healthy volunteers:

- Pregnancy or lactating
- Any medication except contraceptives
- Known underlying diseases: cancer, immune deficiency, hypertension, complement, thrombosis disorder.

## Study design

### Design

Study type:	Observational invasive
Intervention model:	Other
Allocation:	Non-randomized controlled trial
Masking:	Open (masking not used)
Control:	Active
Primary purpose:	Diagnostic

### Recruitment

NL	
Recruitment status:	Recruiting
Start date (anticipated):	13-11-2023
Enrollment:	20
Type:	Actual

## Ethics review

Approved WMO	
Date:	14-04-2023
Application type:	First submission
Review commission:	METC Amsterdam UMC

## Study registrations

### Followed up by the following (possibly more current) registration

No registrations found.

### Other (possibly less up-to-date) registrations in this register

No registrations found.

## In other registers

### Register

CCMO

### ID

NL77890.029.21