ULTRA-ERAS: discharge after 24hours after colorectal surgery, a pilot study

Published: 19-01-2023 Last updated: 14-03-2025

The aim of the study is to assess if hospitalization can be reduced with the use of an early discharge protocol, named *ULTRA-ERAS*, in combination with tele monitoring in patients in elective colorectal surgery

Ethical reviewApproved WMOStatusCompletedHealth condition typeOther conditionStudy typeInterventional

Summary

ID

NL-OMON56276

Source

ToetsingOnline

Brief titleULTRA-ERAS

Condition

- Other condition
- Gastrointestinal neoplasms malignant and unspecified
- Gastrointestinal therapeutic procedures

Synonym

darmaandoening; tumoren/diverticulitis

Health condition

postoperatieve zorg

Research involving

Human

Sponsors and support

Primary sponsor: Sint Antonius Ziekenhuis

Source(s) of monetary or material Support: Onderzoeksfonds Antonius ziekenhuis

Intervention

Keyword: Colorectal, discharge, ERAS

Outcome measures

Primary outcome

Clavien Dindo >= 3 complication rates (%) and initial length of stay (days)

Secondary outcome

success rate of *ULTRA-ERAS* (defined as <24hr length of stay), Total 30d LOS (incl. readmission), 30d complication rate, readmission rates (%), 30d mortality (%), patient satisfaction two weeks after hospitalization measured by 5-likert scale questionnaire, employee satisfaction halfway and after last inclusion of pilot study measured by 5-likert scale questionnaire and semi-structured interviews

Study description

Background summary

Over the last decade, Enhanced Recovery After Surgery (ERAS) protocols are widely integrated in abdominal surgery. This improved the procedures from a medical, financial and logistical perspective. Procedures such as cholecystectomy and bariatric surgery have already been shown to be successfully performed in combination with discharge within twenty-four hours. Current literature suggests that length of stay after colorectal surgery could safely be further reduced, in carefully selected patients by following a clear and structured protocol. Reduction of length of stay might help reduce hospital costs and could increase bed capacity in future healthcare.

Study objective

2 - ULTRA-ERAS: discharge after 24hours after colorectal surgery, a pilot study 3-05-2025

The aim of the study is to assess if hospitalization can be reduced with the use of an early discharge protocol, named *ULTRA-ERAS*, in combination with tele monitoring in patients in elective colorectal surgery

Study design

Prospective pilot intervention study

Intervention

All included patients will be treated according to our ULTRA-ERAS protocol. This protocol aims for discharge within twenty-four hours. After discharge they will receive a telemonitoring app to monitor complaints and vital parameters for the first five days at home.

Study burden and risks

Participants will undergo colorectal surgery and follow the ULTRA-ERAS protocol, which aims at discharge <24 hours after surgery. Previous research has shown that <24hr discharge is safe compared to usual care (1-3). Although patients will be discharge approximately two to three days earlier than usual, we minimize burden and risk by including many checkpoints in our protocol during hospitalization, thereby ensuring patient safety. Patients risk after discharge will be minimized by ensuring a 24/7 telephonic availability in-hospital nurse consultation and digital registration of complaints and vital parameters trough a mobile app.

Contacts

Public

Sint Antonius Ziekenhuis

Koekoekslaan 1 Nieuwegein 3435CM NL

Scientific

Sint Antonius Ziekenhuis

Koekoekslaan 1 Nieuwegein 3435CM NL

Trial sites

Listed location countries

Netherlands

Eligibility criteria

Age

Adults (18-64 years) Elderly (65 years and older)

Inclusion criteria

- Elective minimal invasive right/left/hemicolectomy, ileocoecal or sigmoidal resection
- Uncomplicated procedure at sign-out
- Both benign and malignant indications for surgery
- Age <=80 years
- ASA score I or II
- Complete understanding of procedure and compliance
- A person at home during the first four days and living in a non-public living space (such as Leger des Heils)
- Lives within a half hour travel radius to the hospital and owns transportation to hospital

Exclusion criteria

- Anti-coagulants which require bridging
- Insulin dependent diabetes
- Multi-visceral resections
- Perioperative adhesiolysis, presence of abscess or need for enterostomy
- Perioperative conversion
- Perioperative placement of drains or gastric tubes
- cT4 tumours
- Lives in a public living space (such as Leger des Heils)

Study design

Design

Study type: Interventional

Masking: Open (masking not used)

Control: Uncontrolled

Primary purpose: Treatment

Recruitment

NL

Recruitment status: Completed
Start date (anticipated): 21-03-2023

Enrollment: 30

Type: Actual

Ethics review

Approved WMO

Date: 19-01-2023

Application type: First submission

Review commission: MEC-U: Medical Research Ethics Committees United

(Nieuwegein)

Approved WMO

Date: 05-09-2023

Application type: Amendment

Review commission: MEC-U: Medical Research Ethics Committees United

(Nieuwegein)

Approved WMO

Date: 14-09-2023

Application type: Amendment

Review commission: MEC-U: Medical Research Ethics Committees United

(Nieuwegein)

Approved WMO

Date: 19-10-2023

Application type: Amendment

Review commission: MEC-U: Medical Research Ethics Committees United

(Nieuwegein)

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register ID

CCMO NL81715.100.22