Randomized trial Evaluating first time shoulder Dislocation: sUrgery vs Conservative carE (REDUCE)

Published: 07-06-2024 Last updated: 07-06-2025

The primary objective of the pilot study is to assess the feasibility of a definitive trial to determine the effect of arthroscopic soft tissue stabilization vs. non-surgical treatment on rates of recurrent anterior ...

Ethical review	Approved WMO
Status	Pending
Health condition type	Joint disorders
Study type	Interventional research previously applied in human subjects

Summary

ID

NL-OMON57132

Source ToetsingOnline

Brief title REDUCE

Condition

• Joint disorders

Synonym Shoulder dislocation

Research involving Human

Sponsors and support

Primary sponsor: McMaster University

Source(s) of monetary or material Support: Eerste geldstroom (geld van Ministerie van OC&W aan universiteiten)

1 - Randomized trial Evaluating first time shoulder Dislocation: sUrgery vs Conserva ... 18-06-2025

Intervention

- Surigical procedure
- Other intervention

Keyword: Instability, Shoulder dislocation

Explanation

N.a.

Outcome measures

Primary outcome

The success of the pilot study will be based upon the following a priori
 thresholds:

1. 50 patients recruited within 10 months; < br />

- 2. 42 of 50 participants (85%) achieving complete follow-up at two years; < br />
- 3. 85% of patients allocated to surgical intervention receiving surgery within

- 3 months of enrollment;

- 4. Less than 5 crossovers* between both groups.

*Note: Crossovers are defined as either: 1) participants randomized to

non-operative management (control group) who choose to undergo surgery in the

absence of any feelings of recurrent instability or re-dislocation; or 2)

participants randomized to surgical intervention who opt out of surgery prior

to surgical intervention and choose to undergo primary non-operative management.

We will confirm feasibility with a *traffic light* approach to determine if the

definitive trial will be feasible, require modifications or will not be

feasible.

Secondary outcome

The secondary objectives of the pilot trial will be the clinical objectives of
 the definitive trial:

- 1. Rates of recurrent shoulder dislocations up to 24 months* post-treatment;

- 2. Symptoms of instability without dislocation up to 24 months post treatment; < br />

3. Clinical outcomes measured by Western Ontario Shoulder Instability (WOSI)

Index, American Shoulder and Elbow Society (ASES) score, Shoulder Activity

Scale, EQ-5D, Visual Analog Scale (VAS) Pain Score, and Patient Satisfaction

pr/>
questionnaire;

N

- 4. Physical examination: range of motion, strength, stability;

- 5. Return to previous level of activity and work, and;

6. Safety, shoulder-related complications and serious adverse events.

Hypothesis: We believe that the pilot trial will be feasible in our ability to

Study description

Background summary

The shoulder is the most commonly dislocated joint in the body with a global incidence that ranges from 15 to 25 per 100 000 people. The estimated annual societal cost in North America due to first-time shoulder dislocations exceeds \$1.2 billion CAD.

Anterior dislocations, the most common type of shoulder dislocation, are often complicated by subsequent instability and recurrent dislocation, with reported rates as high as 47%. Shoulder instability commonly results in pain and negatively impacts quality of life.

Current standard of care suggests surgical stabilization of the shoulder after two or more dislocations, but the evidence is far from conclusive. Observational studies suggest that early surgical stabilization has strong biological rationale in limited risks of recurrent dislocation, improving quality of life, and potentially decreasing the future risk of shoulder arthritis. Also, some economic health studies suggests that surgery is less costly and more effective, even after recurrent dislocations.

Study objective

The primary objective of the pilot study is to assess the feasibility of a definitive trial to determine the effect of arthroscopic soft tissue stabilization vs. non-surgical treatment on rates of recurrent anterior dislocation and functional outcomes in patients presenting with a first-time dislocation (FTD) over a 24-month period.

Since the decision between surgery and conservative management may be heavily influenced by patient preference, we will also embed a prospective non-randomized cohort within this RCT to capture all patients who would be eligible for the study but refuse to be randomized. This study design not only ensures maximal participation but is also more generalizable to the real world where preferences might play a role in shared decisionmaking.

Study design

We propose a multi-centre pilot RCT of 50 participants with embedded cohort of an additional 50 participants, across Canada, the United States, South America, and Europe to compare the effect of arthroscopic

soft tissue stabilization (Bankart procedure) and non-operative treatment (physical therapy) in patients with a post-traumatic anterior FTD. Eligible and

3 - Randomized trial Evaluating first time shoulder Dislocation: sUrgery vs Conserva ... 18-06-2025

consenting participants will be followed-up by the site for 24 months. Outcomes will be assessed at 6 weeks, 6 months, 12 months, and 24 months post-treatment.

Intervention

Participants will undergo arthroscopic soft tissue stabilization (Bankart procedure) or non-surgical treatment as described below.

Arthroscopic Stabilization (Intervention)

Patients randomized to surgical intervention will be required to undergo surgery within 3 months of enrollment. Advanced imaging will be obtained prior to surgery, including MRI, MR arthrogram or preoperative CT scan for assessment of pathology. For arthroscopic Bankart repair, the participant will be placed in the lateral decubitus or beach chair position. Standard diagnostic arthroscopy will be performed. The capsulolabral complex will be freed from the anterior aspect of the scapular neck. The anterior aspect of the scapular neck will be decorticated using a motorized burr. A capsuloligamentous repair will be performed with the capsule shifted from inferior to superior and repaired on the glenoid face. The number of anchors used for the repair will be left to the discretion of the surgeon (a minimum of 3 anchors recommended). Remplissage procedure will be performed at the discretion of the operating surgeon. All participants will follow a standardized rehabilitation protocol following surgical intervention: Phase I: Immediate postop phase (0-6 weeks) in sling, Passive range of movement, Phase II: Active motion phase (5-8 weeks after surgery), Phase III: Strengthening phase (8-12 weeks after surgery), Phase IV: Advanced strengthening phase (12 weeks and beyond).

Non-surgical treatment (Control Group)

Type of sling and immobilization will be left to the discretion of the operating surgeon. Patients randomized to non-surgical treatment will begin early physical therapy (PT) immediately as per protocol (2 weeks after initial immobilization). All participants will follow a standardized 3 phase rehabilitation protocol following 2 weeks of immobilization.

Study burden and risks

Following the initial urgent management and relocation of a dislocated shoulder, the prevention of recurrent instability is the critical management consideration for health care providers and surgeons. Two initial management options exist in patients with a FTD: non-operative care or surgical stabilization.

Recent research and available evidence over the past 10 years have called into question the role of a delayed approach to managing FTDs for a number of reasons. Surgical management has been suggested as a more reliable option to prevent further dislocations and improve patient outcomes when compared with non-operative management. Arthroscopic soft tissue repair (Bankart repair) has

become increasingly popular given advancements in surgical technique allowing for a minimally invasive and reliable improvement in shoulder stability with a low risk of complication. The high recurrence rate in younger patients may justify offering surgical treatment after the first dislocation episode. A recent systematic review by Hurley et al.16 found arthroscopic Bankart repair resulted in a 7-fold lower recurrence rate and a higher rate of return to sport and activity than non-operative management21. While other surgical stabilization options exist, including non-anatomic bony transfer (Latarjet procedure), the Bankart repair is widespread as it is minimally invasive and restores native anatomy. Recent data also suggests that patients who are surgically treated following a FTD have improved outcomes when compared to those who have recurrent instability events before undergoing surgery. Delayed management of shoulder instability results in further injury to the shoulder joint. MRI evaluation of individuals who were assessed greater than 6 months from the time of initial dislocation had increased prevalence of not only recurrent shoulder instability events but a greater incidence and severity of intra-articular injury, including SLAP tears, labral tears, and glenoid cartilage damage.

Contacts

Scientific

OLVG S.N.W. Vorrink oosterpark 9 amsterdam 1091 AC Netherlands 020 599 2415 **Public** OLVG S.N.W. Vorrink oosterpark 9 amsterdam 1091 AC Netherlands 020 599 2415

Trial sites

Trial sites in the Netherlands

Deventer Ziekenhuis Target size:

5 - Randomized trial Evaluating first time shoulder Dislocation: sUrgery vs Conserva ... 18-06-2025

5

OLVG	
Target size:	5
Medische Kliniek Velsen	
Target size:	5

Listed location countries

Canada, United States, Kuwait, Spain, Netherlands

Eligibility criteria

Age Adults (18-64 years)

Inclusion criteria

1. Patients ages 18-40 years;

2. Diagnosis of first-time anterior shoulder dislocation having occurred within the past 3 months, confirmed either by radiographic evidence or documented reduction of anterior shoulder dislocation as well as physical examination eliciting unwanted glenohumeral translation with reproduction of symptoms; 3. Provision of informed consent.

Exclusion criteria

- 1. Patients that cannot undergo surgery or anesthesia;
- 2. Patients with concomitant injuries (rotator cuff tear, fracture);
- 3. Previous shoulder surgery;
- 4. Patients that will likely have problems with maintaining follow-up or are incarcerated;
- 5. Epilepsy/seizure disorder
- 6. Pregnancy;
- 7. Diagnosis of multidirectional instability;
- 8. Bony glenoid defect (bony Bankart) >10% as measured on preop imaging;

9. Dislocation without trauma, in a context of hyper laxity or atraumatic instability;

10. Cases involving litigation or workplace insurance claims (e.g. WSIB).

Study design

Design

Study phase:	N/A
Study type:	Interventional research previously applied in human subjects
Intervention model:	Parallel
Allocation:	Randomized controlled trial
Masking:	Open (masking not used)
Control:	Other type of control
Primary purpose:	Treatment

Recruitment

NL	
Recruitment status:	Pending
Start date (anticipated):	02-03-2025
Enrollment:	15
Duration:	24 months (per patient)
Туре:	Anticipated
WORLD	
Recruitment status:	Pending
Start date (anticipated):	01-01-2025
Enrollment:	100
Туре:	Anticipated

Medical products/devices used

Product type:	N.a.
Registration:	No

IPD sharing statement

Plan to share IPD: Undecided

Plan description N.a.

Ethics review

Approved WMO

7 - Randomized trial Evaluating first time shoulder Dislocation: sUrgery vs Conserva ... 18-06-2025

Date:	28-11-2024
Application type:	First submission
Review commission:	MEC-U: Medical Research Ethics Committees United (Nieuwegein)
Approved WMO	
Date:	10-04-2025
Application type:	Amendment
Review commission:	MEC-U

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register CCMO Research portal ID NL86808.100.24 NL-005699