

# Measurements for a safe return to sports after ACL reconstruction.

Gepubliceerd: 20-08-2013 Laatst bijgewerkt: 13-12-2022

The aim of this study is to correlate the score on a functional performance test battery to the occurrence of ACL reruptures and contralateral ruptures. We hypothesize that athletes with a limb symmetry index of

<b>Ethische beoordeling</b>	Positief advies
<b>Status</b>	Werving nog niet gestart
<b>Type aandoening</b>	-
<b>Onderzoekstype</b>	Observationeel onderzoek, zonder invasieve metingen

## Samenvatting

### ID

NL-OMON19946

### Bron

NTR

### Verkorte titel

N/A

### Aandoening

Anterior cruciate ligament reconstruction (ACLR)Voorste kruisband (VKB) reconstructie

Anterior cruciate ligament rupture

Voorste kruisband ruptuur

### Ondersteuning

**Primaire sponsor:** Sport Medisch Centrum Papendal, Arnhem

Universitair Medisch Centrum St Radboud, IQ healthcare, Nijmegen

Rijnstate Ziekenhuis, Arnhem

Kliniek ViaSana, Mill

**Overige ondersteuning:** Fund = initiator = sponsor

### Onderzoeksproduct en/of interventie

## **Uitkomstmaten**

### **Primaire uitkomstmaten**

- Number of graft reruptures and contralateral ACL ruptures two years after ACLR, comparison between subgroups based on limb symmetry index score or qualitative performance.<br>
- Number of injuries at lower limb two years after ACLR, comparison between subgroups based on limb symmetry index score or qualitative performance.

## **Toelichting onderzoek**

### **Achtergrond van het onderzoek**

For athletes rehabilitation after anterior cruciate ligament reconstruction (ACLR) could be judged successful if they return to their previous level of sports. Recent research, however, shows that one third of athletes does not return to this sport level within two years. After three or more years this number even increases to 50%. Half of these athletes report their ACL injury as the primary reason for a lower activity level. Moreover, athletes that do return to their previous sport level have a considerable risk of tearing the graft or the contralateral ACL within the first two years; 3-22% of athletes rerupture the reconstructed ligament and 3-24% rupture the contralateral ACL. The risk of a contralateral ACL injury is higher when returning to a high level of pivoting sports. Three recent systematic reviews show that most clinicians do not use objective criteria for determining the moment of return to sports. Furthermore, there are no measurements for quality of movement performed, despite previous studies showed that, besides quantity, also quality of movement can be a causal factor for ACL (re)ruptures. The presence of dynamic knee valgus when landing from a jump and deficits in neuromuscular control of the trunk can predict ACL injury in healthy athletes. Also, the occurrence of dynamic knee valgus and a smaller knee flexion angle during landing are predictive for ACL rerupture. It can be concluded that both components of functional performance are important in rehabilitation after ACLR and prevention of (re)injuries. Nevertheless, there has never been conducted a prospective cohort study with ACLR athletes, to investigate whether an extensive test battery for measuring quantity and quality of movement is able to predict safe return to sports.

### **Doel van het onderzoek**

The aim of this study is to correlate the score on a functional performance test battery to the occurrence of ACL reruptures and contralateral ruptures. We hypothesize that athletes with a limb symmetry index of <100% on strength or hoptests or a bad quality of movement have more risk for graft rerupture or contralateral rupture.

### **Onderzoeksopzet**

Measurements are conducted at the following timepoints:

T0 - preoperative

T1 - at the end of the first postoperative phase

T2 - at the end of the second postoperative phase

T3 - at the end of the third postoperative phase

T4 - 1 year postoperative

T5 - 1 year and 3 months postoperative

T6 - 1 year and 6 months postoperative

T7 - 1 year and 9 months postoperative

T8 - 2 years postoperative

### **Onderzoeksproduct en/of interventie**

The rehabilitation after ACLR will follow the KNGF Evidence Statement (that will be published in the fall of 2013). Measurement of functional performance are according to the Evidence Statement. The measurements include both quantity and quality of functional performance: strength measurements, hoptest battery, video-analysis for quality of movement during drop jump. Before surgery and after finishing rehabilitation, only an online questionnaire is send, with questions about return to sports and injuries.

## **Contactpersonen**

### **Publiek**

Sport Medisch Centrum Papendal  
Papendallaan 7  
Nicky Engelen-van Melick  
Arnhem 6816 VD  
The Netherlands  
088-088-1350

### **Wetenschappelijk**

Sport Medisch Centrum Papendal  
Papendallaan 7

Nicky Engelen-van Melick  
Arnhem 6816 VD  
The Netherlands  
088-088-1350

## Deelname eisen

### Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

- ACLR from September 2013 until December 2014
- 16-40 years old, male and female
- Athletes with preoperative Tegner Activity scale  $\geq 6$  (only pivoting sports)
- Any arthroscopic surgical method for ACLR with bone-patellar tendon-bone or semitendinosus/gracilis autograft
- All fixation methods
- Concomitant ligament injury (MCL, LCL, PCL, posterolateral corner) grade A or B (IKDC criteria)
- Concomitant meniscectomy
- Cartilage pathology grade I or II (ICRS criteria)
- Patient is willing to rehabilitate at a physiotherapy practice that is a member of the hospitals network
- Patients willing to sign informed consent

### Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

- Autograft or synthetic graft
- Revision surgery
- Meniscal repair

- Concomitant ligament injury C of D
- Cartilage pathology grade III or IV

## Onderzoeksopzet

### Opzet

Type:	Observationeel onderzoek, zonder invasieve metingen
Onderzoeksmodel:	Parallel
Toewijzing:	N.v.t. / één studie arm
Blinding:	Open / niet geblindeerd
Controle:	N.v.t. / onbekend

### Deelname

Nederland	
Status:	Werving nog niet gestart
(Verwachte) startdatum:	01-10-2014
Aantal proefpersonen:	150
Type:	Verwachte startdatum

## Ethische beoordeling

Positief advies	
Datum:	20-08-2013
Soort:	Eerste indiening

## Registraties

### Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

## Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

## In overige registers

Register	ID
NTR-new	NL3845
NTR-old	NTR4130
Ander register	: N/A
ISRCTN	ISRCTN wordt niet meer aangevraagd.

## Resultaten

### Samenvatting resultaten

Intention of publication of study protocol within one year time and of study results after the study is finished.