

Hyperoxic Exercise in Type 2 Diabetes.

Gepubliceerd: 27-04-2010 Laatste bijgewerkt: 13-12-2022

Type 2 diabetes patients frequently suffer from physical deconditioning and vascular disease. Although the exact mechanisms are unclear, microvascular changes and endothelial dysfunction appear to inhibit the oxygen transport and uptake in...

Ethische beoordeling	Positief advies
Status	Deelnemers gezocht
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON20199

Bron

NTR

Verkorte titel

Hyperoxic Exercise in Type 2 Diabetes

Aandoening

Diabetes type 2

Ondersteuning

Primaire sponsor: Erasmus MC, University Medical Center Rotterdam

Overige ondersteuning: Diabetes Fonds

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

The relative improvement in whole-body insulin sensitivity. An insulin sensitivity index (SI) will be calculated by using the minimal model (Bergman et al 1985); a higher SI indicates enhanced insulin sensitivity. Acute-phase insulin

secretion (AIRG) and glucose effectiveness (SG) were also determined from the IVGTT (Bergman et al 1985).

Toelichting onderzoek

Achtergrond van het onderzoek

Rationale:

Type 2 diabetes patients frequently suffer from physical deconditioning and vascular disease. Although the exact mechanisms are unclear, microvascular changes and endothelial dysfunction appear to inhibit the oxygen transport and uptake in peripheral skeletal muscle of type 2 diabetes patients. Hyperoxic-training has been shown a safe intervention to increase exercise capacity and energy expenditure and is now routinely used for athletes ('live-high, train-low') and COPD patients. Furthermore, in COPD patients normalisation of oxygen saturation improves insulin sensitivity.

Objective:

This research project aims to investigate (1) whether hyperoxic interval training improves insulin resistance, endothelial dysfunction, cardiovascular fitness, lipid metabolism and glycemic control in the treatment of deconditioned non-insulin dependent type 2 diabetes patients (2) the pathophysiological role of endothelial dysfunction and insulin resistance on the adaptive response to exercise.

Study design:

Substudy 1 (dose-finding study): intervention study;

Substudy 2 (training intervention study): double-blind placebo controlled intervention study.

Study population:

48 deconditioned (non-)insulin dependent type 2 diabetes patients.

Intervention:

After inclusion, 12 patients will participate in the dose-finding study (substudy 1). The remaining 36 subjects will be randomised and blinded to 16 weeks of progressive 30-45 min cycle ergometer interval training, applied 3 times a week while breathing either hyperoxic (100%O₂, 4-8 L/min) or normoxic (21%O₂-79%N₂, 4-8L/min) humidified air (=placebo) through a nasal tube.

Main study parameters/endpoints:

The main study parameter and end-point is the relative improvement in whole-body insulin sensitivity.

Secondary end-points are relative improvements in vascular function parameters using forearm plethysmography, cardiovascular fitness, systemic blood pressure, body composition, lipid metabolism, glycemic control, GLUT-4 expression in the muscle, mitochondrial enzymatic activity and muscle fiber morphologic characteristics.

Nature and extent of the burden and risks associated with participation, benefit and group relatedness:

Before inclusion subjects will undergo a physical examination and X-ECG to exclude cardiovascular disease and assess cardiorespiratory status. A group of 12 subjects will undergo a submaximal exercise test while obtaining arteriovenous blood samples to optimize oxygen flow during exercise (substudy 1). The other 36 participating subjects will be asked to attend a total of 48 supervised exercise sessions of 45 min (substudy 2). To minimize the risk for a hypoglycemic event during the first 2 weeks of the exercise intervention, capillary blood glucose will be measured following exercise. If necessary medication will be adjusted. Independent of oxygen concentration in the air mixture, patients are expected to improve their physical fitness and metabolic control. Beside the direct therapeutic effects, both interventions are expected to improve general health and well-being.

Although unlikely, in theory hyperoxic exercise training might worsen diabetes related retinopathy. Therefore, an experienced ophthalmologist will monitor and stage diabetic retinopathy before and after the 16 weeks training program. Dual energy x-ray absorptiometry will be used to assess changes in body composition. Frequently sampled intravenous and forearm venous occlusion plethysmography following intrabrachial infusions of methacholine and sodium nitroprusside will be applied. A muscle biopsy will be obtained from the non-dominant vastus lateralis muscle after local anesthesia in 24 subjects (12 from the control and 12 from the intervention group). To prevent bleeding complications patients using low molecular weight heparin derivatives and oral anti-coagulants will be excluded. Furthermore a compressive bandage will be applied for 12 hours. The procedure will be performed under aseptic conditions to prevent infectious complications. Participating subjects will visit the clinical research unit 9 times over a 5 months time period, equivalent to a 22 hours time load. Over a period of 20 weeks multiple arterialised and venous blood samples (equivalent to 258 cc) will be drawn through an intravenous/arterial catheter.

Doel van het onderzoek

Type 2 diabetes patients frequently suffer from physical deconditioning and vascular disease. Although the exact mechanisms are unclear, microvascular changes and endothelial dysfunction appear to inhibit the oxygen transport and uptake in peripheral skeletal muscle of type 2 diabetes patients. Hyperoxic-training has been shown a safe intervention to increase exercise capacity and energy expenditure and is now routinely used for athletes ('live-high, train-low') and COPD patients. Furthermore, in COPD patients normalisation of oxygen saturation improves insulin sensitivity.

Onderzoeksopzet

All participants will receive:

1. Interview;
2. Clinical research and blood pressure measurements;
3. Food diary for three days;
4. Carry an activity monitor for seven days (Actigraph accelerationdevice);
5. Exercise ECG during cycle ergometer interval training;
6. Whole body DEXA measurement;
7. Fundus photography;
8. Frequently sampled intravenous glucose tolerance during three hours;

Onderzoeksproduct en/of interventie

After inclusion, 12 patients will participate in the dosefinding study (substudy 1). The remaining 36 subjects will be randomised and blinded to 16 weeks of progressive 30-45 min cycle ergometer interval training, applied 3 times a week while breathing either hyperoxic (100%O₂, 10-15 L/min) or normoxic (21%O₂-79%N₂, 10-15 L/min) humidified air (=placebo) through mask.

Contactpersonen

Algemeen / deelnemers

P.O. box 2040
D. Visser
Rotterdam 3000 CA
The Netherlands
+31 (0)10 7031679

Wetenschappers

P.O. box 2040
D. Visser
Rotterdam 3000 CA
The Netherlands
+31 (0)10 7031679

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

1. Type 2 diabetes according to WHO criteria for over 2 years;
2. VO₂peak 60-90% of age-predicted value as measured on a cycle-ergometer;
3. Motivated and willing/able to travel 3 times a week to ErasmusMC and participate in a supervised exercise intervention program.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

1. Use of Beta-blocker therapy, low molecular heparin, exogenous insulin therapy (for exercise intervention only), use of oral anti-coagulans therapy;
2. Decompensatio cordis, angina pectoris, myocardial infarction or positive signs of cardiac ischaemia on the ECG during the incremental exercise test;

3. Orthopaedic impairments that would limit participation in the training program;
4. Co-morbidity such as renal failure or > grade III retinopathy or previous diabetic foot ulcer;
5. Cerebro-vascular disease (CVA), neurological diseases or deficits;
6. A history of glaucoma or high intraocular pressure will be contraindication for dilated fundus reflex photography.

Onderzoeksopzet

Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Parallel
Toewijzing:	Toewijzing op basis van loting
Blinding:	Dubbelblind
Controle:	Nepmedicijn

Deelname

Nederland	
Status:	Deelnemers gezocht
(Verwachte) startdatum:	01-05-2010
Aantal proefpersonen:	48
Type:	Onderzoek is nog niet gestart, dit is de verwachte startdatum

Ethische beoordeling

Positief advies	
Datum:	27-04-2010
Soort:	Eerste beoordeling onderzoek

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL2175
NTR-old	NTR2299
Ander register	MEC / EudraCT : 2009-125 / 2009-011448-20 ;
ISRCTN	ISRCTN wordt niet meer aangevraagd.

Resultaten

Samenvatting resultaten

N/A